

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
13 May 2004 (13.05.2004)

PCT

(10) International Publication Number  
**WO 2004/039807 A1**

- (51) International Patent Classification<sup>7</sup>: **C07D 471/14**, (74) Agent: OGILVY RENAULT; Attn: Murphy, Kevin, P., Suite 1600, 1981 McGill College Avenue, Montreal, Québec H3A 2Y3 (CA).  
A61K 31/4375, A61P 29/00, 11/00
- (21) International Application Number: PCT/CA2003/001658 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: 28 October 2003 (28.10.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/422,443 30 October 2002 (30.10.2002) US (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): MERCK FROSST CANADA & CO. [CA/CA]; 16711 Trans-Canada Highway, Kirkland, Québec H9H 3L1 (CA).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): LEBLANC, Yves [CA/CA]; 16711 Trans-Canada Highway, Kirkland, Québec H9H 3L1 (CA). DUFRESNE, Claude [CA/CA]; 16711 Trans-Canada Highway, Kirkland, Québec H9H 3L1 (CA). ROY, Patrick [CA/CA]; 16711 Trans-Canada Highway, Kirkland, Québec H9H 3L1 (CA).
- Published:  
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

WO 2004/039807 A1

(54) Title: PYRIDOPYRROLIZINE AND PYRIDOINDOLIZINE DERIVATIVES

(57) Abstract: Pyridopyrrolizine and pyridoindolizine derivatives are prostaglandin receptor antagonists useful for the treatment of prostaglandin-mediated diseases such as allergic rhinitis, nasal congestion and asthma.

## TITLE OF THE INVENTION

## PYRIDOPYRROLIZINE AND PYRIDOINDOLIZINE DERIVATIVES

## BACKGROUND OF THE INVENTION

The present invention relates to compounds and methods for treating prostaglandin mediated diseases, and certain pharmaceutical compositions thereof. More particularly, the compounds of the invention are structurally different from steroids, antihistamines or adrenergic agonists, and are antagonists of the nasal and pulmonary congestion effects of D-type prostaglandins.


Two review articles describe the characterization and therapeutic relevance of the prostanoid receptors as well as the most commonly used selective agonists and antagonists: *Eicosanoids: From Biotechnology to Therapeutic Applications*, Folco, Samuelsson, Macclouf, and Velo eds, Plenum Press, New York, 1996, chap. 14, 137-154 and *Journal of Lipid Mediators and Cell Signalling*, 1996, 14, 83-87. An article from T. Tsuru *et al.* published in 1997 in *Journal of Medicinal Chemistry*, vol 40, pp.3504-3507 states that "PGD2 is considered to be an important mediator in various allergic diseases such allergic rhinitis, atopic asthma, allergic conjunctivitis and atopic dermatitis." More recently, an article by Matsuoka *et al.* in *Science* (2000), 287:2013-7, describes PGD2 as being a key mediator in allergic asthma. In addition, patents such as US 4,808,608 refer to prostaglandin antagonists as useful in the treatment of allergic diseases, and explicitly allergic asthma. PGD2 antagonists are described in, for example, European Patent Application 837,052 and PCT Application WO98/25919, as well as WO99/62555.

## SUMMARY OF THE INVENTION

The present invention provides novel compounds which are prostaglandin receptor antagonists; more particularly, they are prostaglandin D2 receptor (DP receptor) antagonists. Compounds of the present invention are useful for the treatment of various prostaglandin-mediated diseases and disorders; accordingly the present invention provides a method for the treatment of prostaglandin-mediated diseases using the novel compounds described herein, as well as pharmaceutical compositions containing them.

## DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to compounds of formula I:



I

and pharmaceutically acceptable salts and hydrates thereof, wherein:

A is selected from C<sub>1-3</sub>alkyl optionally substituted with one to four halogen atoms, O(CH<sub>2</sub>)<sub>1-2</sub>, and S(CH<sub>2</sub>)<sub>1-2</sub>;

Ar is aryl or heteroaryl each optionally substituted with one to four groups independently selected from R<sub>g</sub>;

Q is selected from:

- (1) COOH,
- (2) CONR<sup>a</sup>R<sup>b</sup>,
- (3) C(O)NHSO<sub>2</sub>R<sup>c</sup>,
- (4) SO<sub>2</sub>NHR<sup>a</sup>,
- (5) SO<sub>3</sub>H,
- (6) PO<sub>3</sub>H<sub>2</sub>, and
- (7) tetrazolyl;

one of X<sup>1</sup>, X<sup>2</sup>, X<sup>3</sup> or X<sup>4</sup> is nitrogen and the others are independently selected from CH and C-R<sub>g</sub>;

Y<sup>1</sup> is selected from -(CR<sup>d</sup>Re)<sub>a</sub>-X-(CR<sup>d</sup>Re)<sub>b</sub>-, phenylene, C<sub>3-6</sub>cycloalkylidene and C<sub>3-6</sub>cycloalkylene, wherein a and b are integers 0-1 such that the sum of a and b equals 0, 1 or 2, and X is a bond, O, S, NR<sup>a</sup>, C(O), CH(OR<sup>a</sup>), OC(O), C(O)O, C(O)NR<sup>a</sup>, OC(O)NR<sup>a</sup>, NR<sup>a</sup>C(O), CR<sup>d</sup>=CR<sup>e</sup> or C≡C;

Y<sup>2</sup> is selected from (CR<sup>d</sup>Re)<sub>m</sub> and CR<sup>d</sup>=CR<sup>e</sup>;

R<sup>1</sup> is selected from H, CN, OR<sup>a</sup>, S(O)<sub>n</sub>C<sub>1-6</sub>alkyl and C<sub>1-6</sub>alkyl optionally substituted with one to six groups independently selected from halogen, OR<sup>a</sup> and S(O)<sub>n</sub>C<sub>1-6</sub>alkyl;

R<sup>2</sup> is selected from H and C<sub>1-6</sub>alkyl optionally substituted with one to six halogen; or

R<sup>1</sup> and R<sup>2</sup> together represent an oxo; or

R<sup>1</sup> and R<sup>2</sup> taken together form a 3- or 4- membered ring containing 0 or 1 heteroatom selected from N-R<sup>f</sup>, S, and O optionally substituted with one or two groups selected from F, CF<sub>3</sub> and CH<sub>3</sub>;

R<sup>3</sup> is selected from H and C<sub>1-6</sub>alkyl optionally substituted with one to six groups independently selected from OR<sup>a</sup> and halogen;

R<sup>a</sup> and R<sup>b</sup> are independently selected from H, C<sub>1-10</sub>alkyl, C<sub>2-10</sub>alkenyl, C<sub>2-10</sub>alkynyl, Cy and Cy C<sub>1-10</sub>alkyl, wherein said alkyl, alkenyl, alkynyl and Cy are optionally substituted with one to six substituents independently selected from halogen, amino, carboxy, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, aryl, heteroaryl, aryl C<sub>1-4</sub>alkyl, hydroxy, CF<sub>3</sub>, OC(O)C<sub>1-4</sub>alkyl, OC(O)NR<sup>i</sup>R<sup>j</sup>, and aryloxy; or R<sup>a</sup> and R<sup>b</sup> together with the atom(s) to which they are attached form a heterocyclic ring of 4 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R<sup>f</sup>;

R<sup>c</sup> is selected from C<sub>1-6</sub>alkyl optionally substituted with one to six halogen, aryl and heteroaryl, wherein said aryl and heteroaryl are optionally substituted with one to three groups selected from halogen, OC<sub>1-6</sub>alkyl, O-haloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl and haloC<sub>1-6</sub>alkyl;

R<sup>d</sup> and R<sup>e</sup> are independently H, halogen, aryl, heteroaryl, C<sub>1-6</sub>alkyl or haloC<sub>1-6</sub>alkyl;

R<sup>f</sup> is selected from H, C<sub>1-6</sub>alkyl, haloC<sub>1-6</sub>alkyl, Cy, C(O)C<sub>1-6</sub>alkyl, C(O)haloC<sub>1-6</sub>alkyl, and C(O)-Cy;

R<sub>g</sub> is selected from

- (1) halogen,
- (2) CN,
- (3) C<sub>1-6</sub>alkyl optionally substituted with one to eight groups independently selected from aryl, heteroaryl, halogen, NR<sup>a</sup>R<sup>b</sup>, C(O)R<sup>a</sup>, C(OR<sup>a</sup>)R<sup>a</sup>R<sup>b</sup>, SR<sup>a</sup> and OR<sup>a</sup>, wherein aryl, heteroaryl and alkyl are each optionally substituted with one to six groups independently selected from halogen, CF<sub>3</sub>, and COOH,
- (4) C<sub>2-6</sub>alkenyl optionally substituted with one to six groups independently selected from halogen and OR<sup>a</sup>,
- (5) Cy
- (6) C(O)R<sup>a</sup>,
- (7) C(O)OR<sup>a</sup>,
- (8) CONR<sup>a</sup>R<sup>b</sup>,
- (9) OCONR<sup>a</sup>R<sup>b</sup>,

- (10)  $\text{OC}_{1-6}\text{alkyl}$ , wherein alkyl is optionally substituted with one to six substituents selected from halogen, aryl, heteroaryl, OH and  $\text{OC(O)R}^a$ ,
- (11)  $\text{O-Cy}$ ,
- (12)  $\text{S(O)}_n\text{C}_{1-6}\text{alkyl}$ , wherein alkyl is optionally substituted with one to six substituents selected from halogen, aryl, heteroaryl, OH, and  $\text{OC(O)R}^a$ ,
- (13)  $\text{S(O)}_n\text{-Cy}$ ,
- (14)  $-\text{NR}^a\text{S(O)}_n\text{R}^b$ ,
- (15)  $-\text{NR}^a\text{R}^b$ ,
- (16)  $-\text{NR}^a\text{C(O)R}^b$ ,
- (17)  $-\text{NR}^a\text{C(O)OR}^b$ ,
- (18)  $-\text{NR}^a\text{C(O)NR}^a\text{R}^b$ ,
- (19)  $\text{S(O)}_n\text{NR}^a\text{R}^b$ ,
- (20)  $\text{NO}_2$ ,
- (21)  $\text{C}_{5-8}\text{cycloalkenyl}$ ,

wherein Cy is optionally substituted with one to eight groups independently selected from halogen,  $\text{C(O)R}^a$ ,  $\text{OR}^a$ ,  $\text{C}_{1-3}\text{alkyl}$ , aryl, heteroaryl and  $\text{CF}_3$ ;

$\text{R}^i$  and  $\text{R}^j$  are independently selected from hydrogen,  $\text{C}_{1-10}\text{alkyl}$ , Cy and  $\text{Cy-C}_{1-10}\text{alkyl}$ ; or

$\text{R}^i$  and  $\text{R}^j$  together with the nitrogen atom to which they are attached form a ring of 5 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N- $\text{R}^f$ ;

Cy is selected from heterocyclyl, aryl, and heteroaryl;

m is 1, 2 or 3; and

n is 0, 1 or 2.

The invention also encompasses pharmaceutical compositions containing a compound of formula I, and methods for treatment or prevention of prostaglandin mediated diseases using compounds of formula I.

The invention is described using the following definitions unless otherwise indicated.

The term "halogen" or "halo" includes F, Cl, Br, and I.


The term "alkyl" refers to linear, branched and cyclic and bicyclic structures and combinations thereof, containing the indicated number of atoms. Non-limiting examples of alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, s- and t-butyl, pentyl, hexyl, heptyl, octyl, nonyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, eicosyl, 3,7-diethyl-2,2-dimethyl-4-propylnonyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl,

cycloheptyl, cyclopropylmethyl, cyclopentylethyl, methyl substituted cyclopropyl, ethyl substituted cyclobutyl, adamantyl, cyclododecylmethyl, 2-ethyl-1-bicyclo[4.4.0]decyl and the like. For example, the term C<sub>1-6</sub>alkyl encompasses acyclic alkyl groups having the indicated number of carbon atoms as well as -C<sub>x</sub>alkyl-C<sub>z</sub>cycloalkyl wherein x is 0 to 3 and z is 3 to 6 with the proviso that x+z = 3 to 6.

“Cycloalkylidene” refers to the following bivalent radical where the points of attachment are on the same carbon atom:



“Cycloalkylene” refers to the following bivalent radical where the points of attachment are on different carbon atoms:



“Phenylene” refers to the following bivalent radical and includes 1,2-phenylene, 1,3-phenylene and 1,4-phenylene:



“Haloalkyl” means an alkyl group as described above wherein one or more hydrogen atoms have been replaced by halogen atoms, with up to complete substitution of all hydrogen atoms with halo groups. C<sub>1-6</sub>haloalkyl, for example, includes -CF<sub>3</sub>, -CH<sub>2</sub>CF<sub>3</sub>, -CF<sub>2</sub>CF<sub>3</sub> and the like.

“Alkoxy” means alkoxy groups of a straight, branched or cyclic configuration having the indicated number of carbon atoms. C<sub>1-6</sub>alkoxy, for example, includes methoxy, ethoxy, propoxy, isopropoxy, and the like.

“Haloalkoxy” means an alkoxy group as described above in which one or more hydrogen atoms have been replaced by halogen atoms, with up to complete substitution of all

hydrogen atoms with halo groups. C<sub>1-6</sub>haloalkoxy, for example, includes -OCF<sub>3</sub>, -OCH<sub>2</sub>CF<sub>3</sub>, -OCF<sub>2</sub>CF<sub>3</sub> and the like.

"Alkenyl" means linear or branched structures and combinations thereof, of the indicated number of carbon atoms, having at least one carbon-to-carbon double bond, wherein hydrogen may be replaced by an additional carbon-to-carbon double bond. C<sub>2-6</sub>alkenyl, for example, includes ethenyl, propenyl, 1-methylethenyl, butenyl and the like.

"Heterocyclyl" refers to a non-aromatic ring having 1 to 4 heteroatoms said ring being isolated or fused to a second ring selected from 3- to 7-membered alicyclic ring containing 0 to 4 heteroatoms, aryl and heteroaryl, wherein said heteroatoms are independently selected from O, N and S. Non-limiting examples of heterocyclyl include oxetanyl, 1,3-dithiacyclopentane, dihydrobenzofuran, and the like.

"Aryl" means a 6-14 membered carbocyclic aromatic ring system comprising 1-3 benzene rings. If two or more aromatic rings are present, then the rings are fused together, so that adjacent rings share a common bond. Examples include phenyl and naphthyl.

The term "heteroaryl" (Het) as used herein represents a 5-10 membered aromatic ring system containing one ring or two fused rings, 1-4 heteroatoms, selected from O, S and N. Het includes, but is not limited to, tetrazolyl, benzothienyl, quinolinyl, benzothiazolyl, furanyl, pyrimidinyl, purinyl, naphthyridinyl, imidazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, oxazolyl, pyrazolyl, pyridyl, pyrrolyl, tetrazinyl, thiazolyl, thiadiazolyl, thienyl, triazinyl, triazolyl, 1*H*-pyrrole-2,5-dionyl, 2-pyrone, 4-pyrone, pyrrolopyridine, furopyridine and thienopyridine.

"Therapeutically effective amount" means that amount of a drug or pharmaceutical agent that will elicit the biological or medical response of a tissue, a system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician.

The term "treatment" or "treating" includes alleviating, ameliorating, relieving or otherwise reducing the signs and symptoms associated with a disease or disorder.

The term "prophylaxis" means preventing or delaying the onset or the progression of a disease or disorder, or the signs and symptoms associated with such disease or disorder.

The term "composition", as in pharmaceutical composition, is intended to encompass a product comprising the active ingredient(s), and the inert ingredient(s) (pharmaceutically acceptable excipients) that make up the carrier, as well as any product which results, directly or indirectly, from combination, complexation or aggregation of any two or more of the ingredients, or from dissociation of one or more of the ingredients, or from other types of reactions or interactions of one or more of the ingredients. Accordingly, the pharmaceutical

compositions of the present invention encompass any composition made by admixing a compound of Formula I, and pharmaceutically acceptable excipients.

For compounds of formula I, examples of A include, but are not limited to, CH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>, CH<sub>2</sub>CH(CH<sub>3</sub>), CH(Cl), CH<sub>2</sub>CF<sub>2</sub>CH<sub>2</sub>, CH(Cl)CH<sub>2</sub>CH(F), OCH<sub>2</sub>, OCH<sub>2</sub>CH<sub>2</sub>, SCH<sub>2</sub> and SCH<sub>2</sub>CH<sub>2</sub>. Examples of Q include, but are not limited to, CO<sub>2</sub>H, CONH<sub>2</sub>, CONHCH<sub>3</sub>, CONHPh, CON(CH<sub>3</sub>)<sub>2</sub>, CON(CH<sub>2</sub>)<sub>4</sub>, CONHSO<sub>2</sub>CH<sub>3</sub>, SO<sub>2</sub>NHPh, tetrazolyl and the like.

Examples of Y<sup>1</sup> include, but are not limited to, CH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>, CH<sub>2</sub>CH(CH<sub>3</sub>), CH(Cl), CH(Ph), CH<sub>2</sub>CH(CF<sub>3</sub>), CF<sub>2</sub>CH<sub>2</sub>, CH(Cl)CH<sub>2</sub>CH(F), OCH<sub>2</sub>, OCH<sub>2</sub>CH<sub>2</sub>, SCH<sub>2</sub>, CH<sub>2</sub>SCH<sub>2</sub>, S, O, C(O), CH<sub>2</sub>C(O), CH<sub>2</sub>C(O)O, CH<sub>2</sub>C(O)OCH<sub>2</sub>, NH, NHC(O), CH<sub>2</sub>NHC(O), CH<sub>2</sub>NHC(O)CH<sub>2</sub>, CH=CH, CH<sub>2</sub>CH=CHCH<sub>2</sub>, CH<sub>2</sub>C≡C, 1,4-phenylene, 1,1-cyclopropylidene, 1,3-cyclohexylene, and the like.

Examples of Ar include, but are not limited to, phenyl, 2-, 3-, 4-chlorophenyl, 2-, 3-, 4-bromophenyl, 2-, 3-, 4-fluorophenyl, 3,4-dichlorophenyl, 2,3-dichlorophenyl, 2,4-dichlorophenyl, 2,5-dichlorophenyl, 2,6-dichlorophenyl, 3,5-dichlorophenyl, 3-chloro-4-fluorophenyl, 2-chloro-4-fluorophenyl, 4-chloro-2-fluorophenyl, 2-cyanophenyl, 4-methylphenyl, 4-isopropylphenyl, 4-trifluoromethylphenyl, biphenyl, naphthyl, 3-methoxyphenyl, 3-carboxyphenyl, 2-carboxamidophenyl, 4-methoxyphenyl, 3-phenoxyphenyl, 4-(4-pyridyl)phenyl, 4-methylsulfonylphenyl, 3-dimethylaminophenyl, 5-tetrazolyl, 1-methyl-5-tetrazolyl, 2-methyl-5-tetrazolyl, 2-benzothienyl, 2-benzofuranyl, 2-indolyl, 2-quinoliny, 7-quinoliny, 2-benzothiazolyl, 2-benzimidazolyl, 1-benzotriazolyl, 2-furanyl, 3-furanyl, 2-imidazolyl, 5-imidazolyl, 5-isoxazolyl, 4-isoxazolyl, 4-isothiazolyl, 1,2,4-oxadiazol-5-yl, 2-oxazolyl, 4-oxazolyl, 4-pyrazolyl, 5-pyrazolyl, 2-pyridyl, 3-pyridyl, 2-pyrazinyl, 5-pyrimidinyl, 2-pyrrolyl, 4-thiazolyl, 1,2,4-thiadiazol-3-yl, 1,2,5-thiadiazol-4-yl, 1,2,3-thiadiazol-4-yl, 1,2,5-oxadiazol-4-yl, 1,2,3-oxadiazol-4-yl, 1,2,4-triazol-5-yl, 1,2,3-triazol-4-yl, 3-thienyl, 1,2,4-triazol-5-yl, pyrrolo-pyridine, furo[3,2-b]pyridin-2-yl, thieno[2,3-b]pyridin-2-yl, 5(H)-2-oxo-4-furanyl, 5(H)-2-oxo-5-furanyl, (1H,4H)-5-oxo-1,2,4-triazol-3-yl, 4-oxo-2-benzopyranyl, and the like.

Examples of Y<sup>2</sup> include, but are not limited to, CH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>, CH(Cl), CCl<sub>2</sub>, CH(CF<sub>3</sub>), CH(Ph), CH<sub>2</sub>CHCl, C(Cl)=CH<sub>2</sub>, C(Cl)=C(Cl), CH=CH, CH=C(CF<sub>3</sub>), and the like.

Examples of X<sup>1</sup>, X<sup>2</sup>, X<sup>3</sup> and X<sup>4</sup> include, but are not limited to, N, CH, C-CH<sub>3</sub>, C-CH(CH<sub>3</sub>)<sub>2</sub>, C-Ph, C-Cl, C-Br, C-F, C-CF<sub>3</sub>, C-C(O)CH<sub>3</sub>, C-C(O)OH, C-C(O)NH<sub>2</sub>, C-C(O)N(CH<sub>2</sub>)<sub>2</sub>O(CH<sub>2</sub>)<sub>2</sub>, C-OCH<sub>3</sub>, C-OCF<sub>3</sub>, C-OPh, C-SCH<sub>3</sub>, C-SOCH<sub>3</sub>, C-SO<sub>2</sub>CH<sub>3</sub>, C-SO<sub>2</sub>Ph, C-NH<sub>2</sub>, C-N(CH<sub>3</sub>)<sub>2</sub>, C-N(CH<sub>3</sub>)C(O)CH<sub>3</sub>, C-N(CH<sub>3</sub>)C(O)OCH<sub>3</sub>, C-NHC(O)NHCH<sub>3</sub>, C-cyclopropyl, C-cyclobutyl, C-cyclopentyl, and the like.



Examples of  $R^1$  include, but are not limited to, hydrogen, cyano,  $CH_3$ ,  $CH_2CH_3$ ,  $CF_3$ ,  $CH_2CH_2Cl$ , cyclopropyl, and the like.

Examples of  $R^2$  include, but are not limited to, hydrogen,  $CH_3$ ,  $CH_2CH_3$ ,  $CF_3$ ,  $CH_2CH_2Cl$ , cyclopropyl, and the like.

Examples of  $R^3$  include, but are not limited to, hydrogen,  $CH_3$ ,  $CH_2CH_3$ ,  $CF_3$ ,  $CH_2CH_2Cl$ ,  $CH_2CH_2OH$ , cyclopropyl, and the like.

In one embodiment of formula I, the moiety A-Q is  $CH_2CO_2H$ .

In a second embodiment of formula I are compounds wherein the moiety  $Y^1-Ar$  is S-aryl or C(O)-aryl, wherein said aryl is naphthyl or phenyl optionally substituted with 1 to 2 groups selected from  $R^6$ . In one subset thereof,  $Y^1-Ar$  is S-phenyl optionally substituted with 1 or 2 groups selected from halogen,  $C_{1-6}$  alkyl and trifluoromethyl.

In a third embodiment of formula I are compounds wherein  $X^1$  is nitrogen and  $X^2$ ,  $X^3$  and  $X^4$  are independently selected from CH and  $CR^6$ . In one subset, one of  $X^2$ ,  $X^3$  and  $X^4$  is  $CR^6$ , and the others are CH. In another subset one of  $X^2$ ,  $X^3$  and  $X^4$  is CH, and the others are  $CR^6$ .


In a fourth embodiment of formula I are compounds wherein  $X^3$  is nitrogen and  $X^1$ ,  $X^2$  and  $X^4$  are independently selected from CH and  $CR^6$ . In one subset, one of  $X^1$ ,  $X^2$  and  $X^4$  is  $CR^6$ , and the others are CH. In another subset one of  $X^1$ ,  $X^2$  and  $X^4$  is CH, and the others are  $CR^6$ .

In fifth embodiment of formula I are compounds wherein one of  $X^1$ ,  $X^2$  or  $X^3$  is nitrogen and the others are CH or  $CR^6$ , and  $X^4$  is  $CR^6$ . In one subset, one of  $X^1$ ,  $X^2$  or  $X^3$  is nitrogen and the others are CH or  $C-C_{1-6}$ alkyl, and  $X^4$  is  $C-S(O)_n-C_{1-6}$ alkyl or  $C-C_{1-6}$ alkyl optionally substituted with  $OR^a$ .

In sixth embodiment of formula I are compounds wherein  $Y^2$  is selected from  $CH_2$  and  $CH_2CH_2$ .

In seventh embodiment of formula I are compounds where  $R^1$ ,  $R^2$  and  $R^3$  are each hydrogen, or  $R^1$  and  $R^2$  together is oxo, and  $R^3$  is hydrogen.


One group of compounds within formula I is represented by the formula Ia:



Ia

wherein  $X^2$  and  $X^3$  are independently CH or C-Rg, A, Ar, Q,  $Y^1$ ,  $R^1$ ,  $R^2$ , m and Rg are as defined under formula I. In one embodiment of formula Ia are compounds wherein  $X^2$  and  $X^3$  are each CH. In another embodiment are compounds wherein  $R^1$  and  $R^2$  are each H. In yet another embodiment, A-Q is  $CH_2CO_2H$ . In yet another embodiment  $Y^1$ -Ar is S-phenyl optionally substituted with 1 or 2 groups independently selected from halogen, C<sub>1-6</sub> alkyl and trifluoromethyl. In yet another embodiment, Rg is selected from  $SO_2$ -C<sub>1-6</sub>alkyl and C<sub>1-6</sub>alkyl.


Another group of compounds within formula I is represented by the formula Ib:



Ib

wherein  $X^1$  and  $X^2$  are independently CH or C-Rg, A, Ar, Q,  $Y^1$ ,  $R^1$ ,  $R^2$ , m and Rg are as defined under formula I. In one embodiment of formula Ia are compounds wherein  $X^1$  and  $X^2$  are each CH. In another embodiment are compounds wherein  $R^1$  and  $R^2$  are each H. In yet another embodiment, A-Q is  $CH_2CO_2H$ . In yet another embodiment  $Y^1$ -Ar is S-phenyl optionally substituted with 1 or 2 groups independently selected from halogen, C<sub>1-6</sub> alkyl and trifluoromethyl. In yet another embodiment, Rg is selected from  $SO_2$ -C<sub>1-6</sub>alkyl and C<sub>1-6</sub>alkyl.


Another group of compounds within formula I is represented by the formula Ic:



Ic

wherein one of  $X^1$ ,  $X^2$  and  $X^3$  is N and the others are each CH,  $X^4$  is CRG, m is 1 or 2, and Ar,  $Y^1$  and m are as defined under formula I. In one embodiment, Ar is phenyl optionally substituted with 1 or 2 groups independently selected from halogen, C<sub>1</sub>-3alkyl and trifluoromethyl. In another embodiment  $Y^1$  is S or C(O). In yet another embodiment,  $X^4$  is selected from C-S(O)<sub>n</sub>-C<sub>1-6</sub>alkyl and C-C<sub>1-6</sub>alkyl optionally substituted with OR<sup>a</sup>.


Representative compounds of formula I are shown in the following Tables:



X <sup>1</sup>	X <sup>2</sup>	X <sup>3</sup>	X <sup>4</sup>	Ar	Y <sup>1</sup>	m
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	4-Cl-Ph	S	2
N	CH	CH	C(SCH <sub>3</sub> )	4-Cl-Ph	S	2
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	3,4-diCl-Ph	S	2
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	4-Cl-Ph	C(O)	2
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	4-Br-Ph	S	2
CH	CH	N	C(SO <sub>2</sub> CH <sub>3</sub> )	3,4-diCl-Ph	S	1
CH	CH	N	C(SO <sub>2</sub> CH <sub>3</sub> )	3,4-diCl-Ph	S	2
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	4-CF <sub>3</sub> -Ph	S	2
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	2-Cl-4-F-Ph	S	2
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	2-naphthyl	S	2
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	2,3-diCl-Ph	S	2
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	4-CH <sub>3</sub> -Ph	S	2

X1	X2	X3	X4	Ar	Y1	m
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	Ph	S	2
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	2,4-diCl-Ph	S	2
CH	N	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	4-Cl-Ph	S	2
CH	CH	N	C(SO <sub>2</sub> CH <sub>3</sub> )	4-Cl-Ph	S	2
N	C(CH <sub>3</sub> )	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	4-Cl-Ph	S	2
N	CH	C(CH <sub>3</sub> )	C(SO <sub>2</sub> CH <sub>3</sub> )	4-Cl-Ph	S	2
CH	C(CH <sub>3</sub> )	N	C(SO <sub>2</sub> CH <sub>3</sub> )	4-Cl-Ph	S	2
C(CH <sub>3</sub> )	CH	N	C(SO <sub>2</sub> CH <sub>3</sub> )	4-Cl-Ph	S	2
N	CH	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-F-Ph	S	2
N	CH	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-Cl-Ph	S	2
N	CH	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	2,4-diCl-Ph	S	2
N	CH	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-Br-Ph	S	2
N	CH	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	2-Cl-4-F-Ph	S	2
N	CH	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	3,4-diCl-Ph	S	2
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-F-Ph	S	2
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-Cl-Ph	S	2
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	2,4-diCl-Ph	S	2
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-Br-Ph	S	2
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	2-Cl-4-F-Ph	S	2
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	3,4-diCl-Ph	S	2
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-F-Ph	S	1
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-Cl-Ph	S	1
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	2,4-diCl-Ph	S	1
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-Br-Ph	S	1
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	2-Cl-4-F-Ph	S	1
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	3,4-diCl-Ph	S	1
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-F-Ph	S	1
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-Cl-Ph	S	1
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	2,4-diCl-Ph	S	1
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-Br-Ph	S	1
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	2-Cl-4-F-Ph	S	1
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	3,4-diCl-Ph	S	1

X1	X2	X3	X4	Ar	Y1	m
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-F-Ph	S	2
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-Cl-Ph	S	2
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	2,4-diCl-Ph	S	2
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-Br-Ph	S	2
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	2-Cl-4-F-Ph	S	2
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	3,4-diCl-Ph	S	2
N	CH	CH	C(CH(OCH <sub>3</sub> ) (CH <sub>2</sub> CH <sub>3</sub> ))	4-Cl-Ph	S	2
N	CH	CH	C(CH(OCH <sub>3</sub> ) (CH <sub>2</sub> CH <sub>3</sub> ))	4-Cl-Ph	S	1
CH	N	CH	C(CH(OCH <sub>3</sub> ) (CH <sub>2</sub> CH <sub>3</sub> ))	4-Cl-Ph	S	1
CH	N	CH	C(CH(OCH <sub>3</sub> ) (CH <sub>2</sub> CH <sub>3</sub> ))	4-Cl-Ph	S	2
CH	CH	N	C(CH(OCH <sub>3</sub> ) (CH <sub>2</sub> CH <sub>3</sub> ))	4-Cl-Ph	S	2
CH	CH	N	C(CH(OCH <sub>3</sub> ) (CH <sub>2</sub> CH <sub>3</sub> ))	4-Cl-Ph	S	1
N	CH	CH	C(C(CH <sub>3</sub> ) <sub>3</sub> )	4-Cl-Ph	S	2
N	CH	CH	C(C(CH <sub>3</sub> ) <sub>3</sub> )	3,4-diCl-Ph	S	2
N	CH	CH	C(C(CH <sub>3</sub> ) <sub>3</sub> )	4-Br-Ph	S	2
N	CH	CH	C(C(CH <sub>3</sub> ) <sub>3</sub> )	4-CF <sub>3</sub> -Ph	S	2
N	CH	CH	C(C(CH <sub>3</sub> ) <sub>3</sub> )	2-Cl-4-F-Ph	S	2
N	CH	CH	C(C(CH <sub>3</sub> ) <sub>3</sub> )	2-naphthyl	S	2
N	CH	CH	C(C(CH <sub>3</sub> ) <sub>3</sub> )	2,3-diCl-Ph	S	2
N	CH	CH	C(C(CH <sub>3</sub> ) <sub>3</sub> )	4-CH <sub>3</sub> -Ph	S	2
N	CH	CH	C(C(CH <sub>3</sub> ) <sub>3</sub> )	Ph	S	2
N	CH	CH	C(C(CH <sub>3</sub> ) <sub>3</sub> )	2,4-diCl-Ph	S	2



Ar	Y1
5-tetrazolyl	S
2-pyrrolyl	S
1,2,4-triazol-3-yl	S
1,2,3-triazol-4-yl	S
5-imidazolyl	S
4-pyrazolyl	S
5-pyrazolyl	S
(1H,4H)-5-oxo-1,2,4-triazol-3-yl	S
4-isothiazolyl	S
1,2,5-thiadiazol-5-yl	S
1,2,5-oxadiazol-5-yl	S
3-furanyl	S
1,2,3-thiadiazol-4-yl	S
1,2,3-oxadiazol-4-yl	S
4-isoxazolyl	S
3-thienyl	S
4-oxazolyl	S
4-thiazolyl	S
(5H)-2-oxo-5-furanyl	S
(5H)-2-oxo-4-furanyl	S
1,2,4-oxadiazol-5-yl	S
3-pyridyl	S
2-pyrazinyl	S
5-pyrimidinyl	S
2-indolyl	S
2-benzothieryl	S

Ar	Y1
2-benzofuranyl	S
4-oxo-benzopyran-2-yl	S
2-quinolinyl	S
2-benzimidazolyl	S
2-benzoxazolyl	S
2-benzothiazolyl	S
1-benzotriazolyl	CH <sub>2</sub> S
thieno[2,3-b]pyridin-2-yl	S

For purposes of this specification, the following abbreviations have the indicated meanings:

Ac	=	acetyl
AcO	=	acetate
BOC	=	t-butyloxycarbonyl
CBZ	=	carbobenzyloxy
CDI	=	carbonyldiimidazole
DCC	=	1,3-dicyclohexylcarbodiimide
DCE	=	1,2-dichloroethane
DIBAL	=	diisobutyl aluminum hydride
DIEA	=	N,N-diisopropylethylamine
DMAP	=	4-(dimethylamino)pyridine
DMF	=	dimethylformamide
EDCI	=	1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
EDTA	=	ethylenediaminetetraacetic acid, tetrasodium salt hydrate
FAB	=	fast atom bombardment
Fmoc	=	9-fluorenylmethoxycarbonyl
HMPA	=	hexamethylphosphoramide
HATU	=	O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate
HOBt	=	1-hydroxybenzotriazole
HRMS	=	high resolution mass spectrometry

ICBF	=	isobutyl chloroformate
KHMDS	=	potassium hexamethyldisilazane
LDA	=	lithium diisopropylamide
MCPBA	=	metachloroperbenzoic acid
MMPP	=	magnesium monoperoxyphthalate hexahydrate
Ms	=	methanesulfonyl = mesyl
MsO	=	methanesulfonate = mesylate
NBS	=	N-bromosuccinimide
NMM	=	4-methylmorpholine
NMP	=	N-methylpyrrolidinone
PCC	=	pyridinium chlorochromate
PDC	=	pyridinium dichromate
Ph	=	phenyl
PPTS	=	pyridinium p-toluene sulfonate
pTSA	=	p-toluene sulfonic acid
PyH•Br <sub>3</sub>	=	pyridine hydrobromide perbromide
r.t./RT	=	room temperature
rac.	=	racemic
TFA	=	trifluoroacetic acid
TfO	=	trifluoromethanesulfonate = triflate
THF	=	tetrahydrofuran
TLC	=	thin layer chromatography
TMSCl	=	trimethylsilyl chloride

## Alkyl group abbreviations

Me	=	methyl
Et	=	ethyl
n-Pr	=	normal propyl
i-Pr	=	isopropyl
c-Pr	=	cyclopropyl
n-Bu	=	normal butyl
i-Bu	=	isobutyl
c-Bu	=	cyclobutyl
s-Bu	=	secondary butyl
t-Bu	=	tertiary butyl



### Optical Isomers - Diastereomers - Tautomers

Compounds of formula I contain one or more asymmetric centers and can thus occur as racemates and racemic mixtures, single enantiomers, diastereomeric mixtures and individual diastereomers. The present invention is meant to comprehend all such isomeric forms of the compounds of formula I.

Some of the compounds described herein may exist with different points of attachment of hydrogen, referred to as tautomers. Such an example may be a ketone and its enol form known as keto-enol tautomers. The individual tautomers as well as mixture thereof are encompassed with compounds of formula I.

Compounds of formula I may be separated into diastereoisomeric pairs of enantiomers by, for example, fractional crystallization from a suitable solvent, for example methanol or ethyl acetate or a mixture thereof. The pair of enantiomers thus obtained may be separated into individual stereoisomers by conventional means, for example by the use of an optically active acid or base as a resolving agent, or by chiral separation techniques such as separation by HPLC using a chiral column.

Alternatively, any enantiomer of a compound of the general formula I or Ia may be obtained by stereospecific synthesis using optically pure starting materials or reagents of known configuration.

### Salts

The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases including inorganic bases and organic bases. Salts derived from inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic salts, manganous, potassium, sodium, zinc, and the like. Particularly preferred are the ammonium, calcium, magnesium, potassium, and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, and basic ion exchange resins, such as arginine, betaine, caffeine, choline, N,N'-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethyl-morpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine, and the like.

When the compound of the present invention is basic, salts may be prepared from pharmaceutically acceptable non-toxic acids, including inorganic and organic acids. Such acids include acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, pantoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic acid, and the like. Particularly preferred are citric, hydrobromic, hydrochloric, maleic, phosphoric, sulfuric, and tartaric acids.

It will be understood that, unless otherwise specified, references to the compound of formula I are meant to also include the pharmaceutically acceptable salts.

### Utilities

Compounds of formula I are antagonists of prostaglandin D2. The ability of compounds of formula I to interact with prostaglandin D2 receptor makes them useful for preventing or reversing undesirable symptoms caused by prostaglandins in a mammalian, especially human subject. The antagonism of the actions of prostaglandin D2 indicates that the compounds and pharmaceutical compositions thereof are useful to treat, prevent, or ameliorate in mammals and especially in humans: respiratory conditions, allergic conditions, pain, inflammatory conditions, mucus secretion disorders, bone disorders, sleep disorders, fertility disorders, blood coagulation disorders, trouble of the vision as well as immune and autoimmune diseases. In addition, such a compound may inhibit cellular neoplastic transformations and metastatic tumor growth and hence can be used in the treatment of cancer. Compounds of formula I may also be of use in the treatment and/or prevention prostaglandin D2 mediated proliferation disorders such as may occur in diabetic retinopathy and tumor angiogenesis. Compounds of formula I may also inhibit prostanoid-induced smooth muscle contraction by antagonizing contractile prostanoids or mimicking relaxing prostanoids and hence may be use in the treatment of dysmenorrhea, premature labor and eosinophil related disorders.

Accordingly, another aspect of the invention provides a method of treating or preventing a prostaglandin D2 mediated disease comprising administering to a mammalian patient in need of such treatment a compound of formula I in an amount which is effective for treating or preventing said prostaglandin D2 mediated disease. Prostaglandin D2 mediated diseases include, but are not limited to, allergic rhinitis, nasal congestion, rhinorrhea, perennial rhinitis, nasal inflammation, asthma including allergic asthma, chronic obstructive pulmonary diseases and other forms of lung inflammation; pulmonary hypotension; sleep disorders and sleep-wake cycle disorders; prostanoid-induced smooth muscle contraction associated with dysmenorrhea and premature labor; eosinophil related disorders; thrombosis; glaucoma and

vision disorders; occlusive vascular diseases, such as for example atherosclerosis; congestive heart failure; diseases or conditions requiring a treatment of anti-coagulation such as post-injury or post surgery treatment; rheumatoid arthritis and other inflammatory diseases; gangrene; Raynaud's disease; mucus secretion disorders including cytoprotection; pain and migraine; diseases requiring control of bone formation and resorption such as for example osteoporosis; shock; thermal regulation including fever; rejection in organ transplant and by-pass surgery, and immune disorders or conditions in which immunoregulation is desirable. More particularly the disease to be treated is one mediated by prostaglandin D2 such as nasal congestion, allergic rhinitis, pulmonary congestion, and asthma including allergic asthma.

### Dose Ranges

The magnitude of prophylactic or therapeutic dose of a compound of formula I will, of course, vary with the nature and the severity of the condition to be treated and with the particular compound of formula I and its route of administration. It will also vary according to a variety of factors including the age, weight, general health, sex, diet, time of administration, rate of excretion, drug combination and response of the individual patient. In general, the daily dose from about 0.001 mg to about 100 mg per kg body weight of a mammal, preferably 0.01 mg to about 10 mg per kg. On the other hand, it may be necessary to use dosages outside these limits in some cases.

The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. For example, a formulation intended for the oral administration of humans may contain from 0.05 mg to 5 g of active agent compounded with an appropriate and convenient amount of carrier material which may vary from about 5 to about 99.95 percent of the total composition. Dosage unit forms will generally contain between from about 0.1 mg to about 0.4 g of an active ingredient, typically 0.5 mg, 1 mg, 2 mg, 5 mg, 10 mg, 25 mg, 50 mg, 100 mg, 200 mg, or 400 mg.

### Pharmaceutical Compositions

Another aspect of the present invention provides pharmaceutical compositions comprising a compound of formula I with a pharmaceutically acceptable carrier. The term "composition", as in pharmaceutical composition, is intended to encompass a product comprising the active ingredient(s), and the inert ingredient(s) (pharmaceutically acceptable excipients) that make up the carrier, as well as any product which results, directly or indirectly, from combination, complexation or aggregation of any two or more of the ingredients, or from

dissociation of one or more of the ingredients, or from other types of reactions or interactions of one or more of the ingredients. Accordingly, the pharmaceutical compositions of the present invention encompass any composition made by admixing a compound of Formula I, additional active ingredient(s), and pharmaceutically acceptable excipients.

For the treatment of any of the prostanoid mediated diseases compounds of formula I may be administered orally, by inhalation spray, topically, parenterally or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. In addition to the treatment of warm-blooded animals such as mice, rats, horses, cattle, sheep, dogs, cats, etc., the compound of the invention is effective in the treatment of humans.

The pharmaceutical compositions containing the active ingredient may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavouring agents, colouring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example, magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated by the technique described in the U.S. Patent 4,256,108; 4,166,452; and 4,265,874 to form osmotic therapeutic tablets for control release.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredients is mixed with water-miscible solvents such as propylene glycol, PEGs and ethanol, or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions contain the active material in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropyl methylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethylene-oxyctanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl, p-hydroxybenzoate, one or more colouring agents, one or more flavouring agents, and one or more sweetening agents, such as sucrose, saccharin or aspartame.

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavouring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavouring and colouring agents, may also be present.

The pharmaceutical compositions of the invention may also be in the form of an oil-in-water emulsion. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavouring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent,

a preservative and flavouring and colouring agents. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butane diol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. Cosolvents such as ethanol, propylene glycol or polyethylene glycols may also be used. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

Compounds of formula I may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ambient temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

For topical use, creams, ointments, gels, solutions or suspensions, etc., containing the compound of formula I are employed. (For purposes of this application, topical application shall include mouth washes and gargles.) Topical formulations may generally be comprised of a pharmaceutical carrier, cosolvent, emulsifier, penetration enhancer, preservative system, and emollient.

#### Combinations with Other Drugs

For the treatment and prevention of prostaglandin mediated diseases, compound of formula I may be co-administered with other therapeutic agents. Thus in another aspect the present invention provides pharmaceutical compositions for treating prostaglandin D2 mediated diseases comprising a therapeutically effective amount of a compound of formula I and one or more other therapeutic agents. Suitable therapeutic agents for combination therapy with a compound of formula I include: (1) a prostaglandin receptor antagonist; (2) a corticosteroid such as triamcinolone acetonide; (3) a  $\beta$ -agonist such as salmeterol, formoterol, terbutaline, metaproterenol, albuterol and the like; (4) a leukotriene modifier, such as a leukotriene antagonist or a lipooxygenase inhibitor such as montelukast, zafirlukast, pranlukast, or zileuton; (5) an antihistamine (histamine H1 antagonist) such as bromopheniramine, chlorpheniramine, dexchlorpheniramine, triprolidine, clemastine, diphenhydramine, diphenylpyraline, tripeleminamine, hydroxyzine, methdilazine, promethazine, trimeprazine, azatadine,

cyproheptadine, antazoline, pheniramine, pyrilamine, astemizole, norastemizole, terfenadine, loratadine, cetirizine, levocetirizine, fexofenadine, desloratadine, and the like; (6) a decongestant including phenylephrine, phenylpropanolamine, pseudophedrine, oxymetazoline, ephinephrine, naphazoline, xylometazoline, propylhexedrine, or levo-desoxyephedrine; (7) an anti-tussive including codeine, hydrocodone, caramiphen, carbapentane, or dexamethorphan; (8) another prostaglandin ligand including prostaglandin F agonist such as latanoprost; misoprostol, enprostil, rioprostil, ornoprostol or rosaprostol; (9) a diuretic; (10) non-steroidal antiinflammatory agents (NSAIDs) such as propionic acid derivatives (alminoprofen, benoxaprofen, bucloxic acid, carprofen, fenbufen, fenoprofen, fluprofen, flurbiprofen, ibuprofen, indoprofen, ketoprofen, miroprofen, naproxen, oxaprozin, piroprofen, pranoprofen, suprofen, tiaprofenic acid, and tioxaprofen), acetic acid derivatives (indomethacin, acemetacin, alclofenac, clidanac, diclofenac, fenclofenac, fenclozic acid, fentiazac, furofenac, ibufenac, isoxepac, oxpinac, sulindac, tiopinac, tolmetin, zidometacin, and zomepirac), fenamic acid derivatives (flufenamic acid, meclofenamic acid, mefenamic acid, niflumic acid and tolfenamic acid), biphenylcarboxylic acid derivatives (diflunisal and flufenisal), oxicams (isoxicam, piroxicam, sudoxicam and tenoxicam), salicylates (acetyl salicylic acid, sulfasalazine) and the pyrazolones (apazone, bezpiperylon, feprazone, mofebutazone, oxyphenbutazone, phenylbutazone); (11) cyclooxygenase-2 (COX-2) inhibitors such as celecoxib and rofecoxib, etoricoxib and valdecoxib; (12) inhibitors of phosphodiesterase type IV (PDE-IV) e.g. Ariflo, roflumilast; (13) antagonists of the chemokine receptors, especially CCR-1, CCR-2, and CCR-3; (14) cholesterol lowering agents such as HMG-CoA reductase inhibitors (lovastatin, simvastatin and pravastatin, fluvastatin, atorvastatin, and other statins), sequestrants (cholestyramine and colestipol), nicotinic acid, fenofibric acid derivatives (gemfibrozil, clofibrat, fenofibrate and benzaifibrate), and probucol; (15) anti-diabetic agents such as insulin, sulfonylureas, biguanides (metformin),  $\alpha$ -glucosidase inhibitors (acarbose) and glitazones (troglitazone, pioglitazone, englitazone, rosiglitazone and the like); (16) preparations of interferon beta (interferon beta-1a, interferon beta-1b); (17) anticholinergic agents such as muscarinic antagonists (ipratropium bromide and tiotropium bromide), as well as selective muscarinic M3 antagonists; (18) steroids such as beclomethasone, methylprednisolone, betamethasone, prednisone, dexamethasone, and hydrocortisone; (19) triptans commonly used for the treatment of migraine such as sumatriptan and rizatriptan; (20) alendronate and other treatments for osteoporosis; (21) other compounds such as 5-aminosalicylic acid and prodrugs thereof, antimetabolites such as azathioprine and 6-mercaptopurine, cytotoxic cancer chemotherapeutic agents, bradykinin (BK2 or BK1) antagonists, TP receptor antagonists such as seratrodist, neurokinin antagonists (NK1/NK2), VLA-4 antagonists such as those described in US 5,510,332, WO97/03094, WO97/02289,

WO96/40781, WO96/22966, WO96/20216, WO96/01644, WO96/06108, WO95/15973 and WO96/31206.

In addition, the invention encompasses a method of treating prostaglandin D<sub>2</sub> mediated diseases comprising: administering to a patient in need of such treatment a therapeutically effective amount of the compound of formula I, co-administered with one or more of such ingredients as listed immediately above. The amounts of active ingredients may be those commonly used for each active ingredient when it is administered alone, or in some instances the combination of active ingredients may result in lower dosage for one or more of the active ingredients.

## METHODS OF SYNTHESIS

Compounds of Formula I of the present invention may be prepared according to the synthetic routes outlined in Schemes A to F and by following the methods described in the Examples provided herein. Both the schemes and specific examples provided herein are for illustrative purpose, and a person skilled in the art will appreciate that other compounds of the present invention may be analogously prepared using the illustrative procedures, or they may be obtained from exemplified compounds via functional group interconversion procedures that are generally known in the art, or they may be prepared by other procedures that are known to persons skilled in the art of organic synthesis.


### Method A

Pyridine 1 can be formylated to give aldehyde 2 according to the procedure described in *J. Heterocyclic Chem.*, p 81, (1988), *Heterocycles* p. 151, 1993 or in *Synthesis*, p. 306 (1999). Halogen displacement with sodium thiomethoxide or sodium methoxide, followed by condensation with methylazidoacetate provides azido olefin 4 which is cyclized under thermal conditions to give indole 5 (see for example, *Tetrahedron Lett.*, 2000, 41:4777-4780). For the fused five-membered ring series, 5 is treated with methyl acrylate in the presence of K<sup>+</sup>OT<sup>-</sup>Bu, followed by decarboxylation with HCl/EtOH to give 6 (m=1). For the fused six and seven-membered ring series, 5 is treated with NaH/DMF and the appropriate bromo ester, followed by cyclization with K<sup>+</sup>OT<sup>-</sup>Bu/THF and finally HCl/EtOH to effect decarboxylation to give 6 (m=2,3). The formation of ester 7 is carried out using Reformatsky conditions followed by deoxygenation with TMSI/NaI, or via Horner-Emmonds reaction followed by hydrogenation over PtO<sub>2</sub> or Pd(OH)<sub>2</sub>. Alternatively, 6 can be converted to 7 by reduction with NaBH<sub>4</sub> in ethanol-THF followed by reaction with diphenyl chlorophosphate using NaHMDS as a base. The resulting



phosphate is treated with dimethyl malonate and NaHMDS. The bis ester is heated in DMSO with NaCl to provide 7.


Scheme A



## Method B

Friedel-Crafts reaction of 7 with a suitable acid chloride in 1,2-dichloroethane gives the corresponding ketone 8. Subsequent cleavage of the ester is effected with sodium hydroxide to give the acid 9. To prepare the thioether 11, a suitable disulfide is treated with  $\text{SO}_2\text{Cl}_2$  in 1,2-dichloroethane to give the corresponding sulfenyl chloride, which is then allowed to react with 7 to give the thioether ester 10. Hydrolysis of 10 using aqueous sodium hydroxide provides the acid 11.


## Scheme B



## Method C

The thioether compound 8a can be oxidized with  $\text{Na}_2\text{WO}_4/\text{H}_2\text{O}_2$  to afford the corresponding sulfone ester, which upon hydrolysis provides the sulfone acid 12. Compound 7a can be similarly oxidized, and the resulting sulfone 13 can be elaborated to the thioether 14 according to procedures described in Scheme B.

## Scheme C




## Method D

Bromopyridine aldehyde 2a is elaborated into compound 15 using the reaction steps described in Method A. Introduction of the ester and isopropyl moieties is as follows. Reformatsky reaction followed by deoxygenation with  $\text{TMSCl}/\text{NaI}$ , then palladium mediated coupling with 2-bromopropene followed by hydrogenation gives 16. Alternatively, Horner-Emmonds reaction followed by palladium mediated coupling with 2-bromopropene and finally hydrogenation of the two olefins gives 16, which is elaborated into compound 17 as described in Scheme B. The isopropyl group can also be introduced earlier in the synthesis. The compound 2a


can be converted to the azaindole ester as shown in Method A for the preparation of 5. The isopropyl is then introduced as described using palladium mediated coupling with 2-bromopropene.



### Scheme D



- 1)  $\text{Br}-\text{CH}_2-\text{CO}_2\text{Me}$ , Zn/Cu couple /THF
- 2) TMSCl, NaI/ $\text{CH}_3\text{CN}$
- 3)  $\text{dppfPdCl}_2 \cdot \text{CH}_2\text{Cl}_2$ ,  $\text{Mg}$ ,  $\text{ZnBr}_2$ ,  $\text{CH}_2=\text{CHBr}$
- 4)  $\text{H}_2$  /  $\text{PtO}_2$  or  $\text{Pd}(\text{OH})_2$

**or**

- 1)  $\text{Et}_2\text{O}_3\text{P}-\text{CH}_2-\text{CO}_2\text{Et}$ , NaH/DMF
- 2)  $\text{dppfPdCl}_2 \cdot \text{CH}_2\text{Cl}_2$ , ,  
Mg,  $\text{ZnBr}_2$
- 3)  $\text{H}_2$  /  $\text{PtO}_2$  or  $\text{Pd}(\text{OH})_2$




**m=1,2,3**  
**R = Me, Et**


## Method E

Azaindole 18 can be prepared according to the procedure in J. Heterocyclic Chem. 359 (1992). Treatment of 18 with base, followed by CO<sub>2</sub> and diazomethane gives ester 19, which is then further functionalized by chemistry described in Methods A and B to give acid

23. Alternatively, condensation of 2-pyridinecarboxaldehyde with methylazidoacetate provides azido olefin 25 which is cyclized under thermal conditions to give ester 19 which is carried on to acid 23.




alternatively:



#### Method F

Compound 26 is prepared from 2-chloropyridine according to the general method described in Method A. The ketone 26 is then converted into the methylvinyl compound 27 using tributylisopropenylstannane (*J. Org. Chem.* 1988 page 3218) and tris(dibenzylidene acetone)dipalladium in the presence of triphenylarsine. Wittig Horner on ketone 27 followed by hydrogenation provides ester 28. The ester 28 is converted to compound 29. The enantiomers are separated on chiral HPLC OD column followed by hydrolysis to provide 30 and 31



#### ASSAYS FOR DETERMINING BIOLOGICAL ACTIVITY

Compounds of formula I can be tested using the following assays to determine their prostanoid antagonist or agonist activity *in vitro* and *in vivo* and their selectivity. The prostaglandin receptor activities demonstrated are DP, EP<sub>1</sub>, EP<sub>2</sub>, EP<sub>3</sub>, EP<sub>4</sub>, FP, IP and TP.

#### Stable expression of prostanoid receptors in the human embryonic kidney (HEK) 293(ebna) cell line

Prostanoid receptor cDNAs corresponding to full length coding sequences are subcloned into the appropriate sites of mammalian expression vectors and transfected into HEK 293(ebna) cells. HEK 293(ebna) cells expressing the individual cDNAs are grown under selection and individual colonies are isolated after 2-3 weeks of growth using the cloning ring method and subsequently expanded into clonal cell lines.

#### Prostanoid receptor binding assays

HEK 293(ebna) cells are maintained in culture, harvested and membranes are prepared by differential centrifugation, following lysis of the cells in the presence of protease inhibitors, for use in receptor binding assays. Prostanoid receptor binding assays are performed in 10 mM MES/KOH (pH 6.0) (EPs, FP and TP) or 10 mM HEPES/KOH (pH 7.4) (DP and IP), containing 1 mM EDTA, 10 mM divalent cation and the appropriate radioligand. The reaction is initiated by addition of membrane protein. Ligands are added in dimethylsulfoxide which is kept constant at 1 % (v/v) in all incubations. Non-specific binding is determined in the presence of 1  $\mu$ M of the corresponding non-radioactive prostanoid. Incubations are conducted for 60 min at room temperature or 30 °C and terminated by rapid filtration. Specific binding is calculated by subtracting non specific binding from total binding. The residual specific binding at each ligand concentration is calculated and expressed as a function of ligand concentration in order to construct sigmoidal concentration-response curves for determination of ligand affinity.

#### Prostanoid receptor agonist and antagonist assays

Whole cell second messenger assays measuring stimulation (EP<sub>2</sub>, EP<sub>4</sub>, DP and IP in HEK 293(ebna) cells) or inhibition (EP<sub>3</sub> in human erythroleukemia (HEL) cells) of intracellular cAMP accumulation or mobilization of intracellular calcium (EP<sub>1</sub>, FP and TP in HEK 293(ebna) cells stably transfected with apo-aequorin) are performed to determine whether receptor ligands are agonists or antagonists. For cAMP assays, cells are harvested and resuspended in HBSS containing 25 mM HEPES, pH 7.4. Incubations contain 100  $\mu$ M RO-20-1724 (phosphodiesterase type IV inhibitor, available from Biomol) and, in the case of the EP<sub>3</sub> inhibition assay only, 15  $\mu$ M forskolin to stimulate cAMP production. Samples are incubated at 37°C for 10 min, the reaction is terminated and cAMP levels are then measured. For calcium mobilization assays, cells are charged with the co-factors reduced glutathione and coelenterazine, harvested and resuspended in Ham's F12 medium. Calcium mobilization is measured by monitoring luminescence provoked by calcium binding to the intracellular photoprotein aequorin. Ligands are added in dimethylsulfoxide which is kept constant at 1 % (v/v) in all incubations. For agonists, second messenger responses are expressed as a function of ligand concentration and both EC<sub>50</sub> values and the maximum response as compared to a prostanoid standard are calculated. For antagonists, the ability of a ligand to inhibit an agonist response is determined by Schild analysis and both K<sub>B</sub> and slope values are calculated.



Prevention of PGD2 or allergen induced nasal congestion in allergic sheep

Animal preparation: Healthy adult sheeps (18-50 kg) are used. These animals are selected on the basis of a natural positive skin reaction to an intradermal injection of *Ascaris suum* extract.

Measurements of nasal congestion: The experiment is performed on conscious animals. They are restrained in a cart in a prone position with their heads immobilized. Nasal airway resistance (NAR) is measured using a modified mask rhinometry technique. A topical anaesthesia (2% lidocaine) is applied to the nasal passage for the insertion of a nasotracheal tube. The maximal end of the tube is connected to a pneumotachograph and a flow and pressure signal is recorded on an oscilloscope linked to a computer for on-line calculation of NAR. Nasal provocation is performed by the administration of an aerosolized solution (10 puffs/nostril). Changes in the NAR congestion are recorded prior to and for 60-120 minutes post-challenge.

Prevention of PGD2 and allergen induced nasal obstruction in cynomolgus monkey

Animal preparation: Healthy adult male cynomologus monkeys (4-10 kg) are used. These animals are selected on the basis of a natural positive skin reaction to an intradermal injection of *Ascaris suum* extract. Before each experiment, the monkey selected for a study is fasted overnight with water provided *ad libitum*. The next morning, the animal is sedated with ketamine (10-15 mg/kg i.m.) before being removed from its home cage. It is placed on a heated table (36°C) and injected with a bolus dose (5-12 mg/kg i.v.) of propofol. The animal is intubated with a cuffed endotracheal tube (4-6 mm I.D.) and anaesthesia is maintained via a continuous intravenous infusion of propofol (25-30 mg/kg/h). Vital signs (heart rate, blood pressure, respiratory rate, body temperature) are monitored throughout the experiment.

Measurements of nasal congestion: A measurement of the animal respiratory resistance is taken via a pneumotachograph connected to the endotracheal tube to ensure that it is normal. An Ecovision acoustic rhinometer is used to evaluate nasal congestion. This technique gives a non-invasive 2D echogram of the inside of the nose. The nasal volume and the minimal cross-sectional area along the length of the nasal cavity are computed within 10 seconds by a laptop computer equipped with a custom software (Hood Laboratories, Mass, U.S.A.). Nasal challenge is delivered directly to the animal's nasal cavity (50 µL volume). The changes in nasal congestion are recorded prior to and for 60-120 minutes post-challenge. If nasal congestion occurs, it will translate into a reduction in the nasal volume.

### Pulmonary Mechanics in Trained Conscious Squirrel Monkeys

The test procedure involves placing trained squirrel monkeys in chairs in aerosol exposure chambers. For control purposes, pulmonary mechanics measurements of respiratory parameters are recorded for a period of about 30 minutes to establish each monkey's normal control values for that day. For oral administration, compounds are dissolved or suspended in a 1% methocel solution (methylcellulose, 65HG, 400 cps) and given in a volume of 1 mL/kg body weight. For aerosol administration of compounds, a DeVilbiss ultrasonic nebulizer is utilized. Pretreatment periods vary from 5 minutes to 4 hours before the monkeys are challenged with aerosol doses of either PGD2 or *Ascaris suum* antigen; 1:25 dilution.

Following challenge, each minute of data is calculated by computer as a percent change from control values for each respiratory parameter including airway resistance ( $R_L$ ) and dynamic compliance ( $C_{dyn}$ ). The results for each test compound are subsequently obtained for a minimum period of 60 minutes post challenge which are then compared to previously obtained historical baseline control values for that monkey. In addition, the overall values for 60 minutes post-challenge for each monkey (historical baseline values and test values) are averaged separately and are used to calculate the overall percent inhibition of mediator or *Ascaris* antigen response by the test compound. For statistical analysis, paired t-test is used. (References: McFarlane, C.S. et al., Prostaglandins, 28, 173-182 (1984) and McFarlane, C.S. et al., Agents Actions, 22, 63-68 (1987).)

### Prevention of Induced Bronchoconstriction in Allergic Sheep

**Animal Preparation:** Adult sheep with a mean weight of 35 kg (range, 18 to 50 kg) are used. All animals used meet two criteria: a) they have a natural cutaneous reaction to 1:1,000 or 1:10,000 dilutions of *Ascaris suum* extract (Greer Diagnostics, Lenois, NC); and b) they have previously responded to inhalation challenge with *Ascaris suum* with both an acute bronchoconstriction and a late bronchial obstruction (W.M. Abraham et al., Am. Rev. Resp. Dis., 128, 839-44 (1983)).

**Measurement of Airway Mechanics:** The unsedated sheep are restrained in a cart in the prone position with their heads immobilized. After topical anesthesia of the nasal passages with 2% lidocaine solution, a balloon catheter is advanced through one nostril into the lower esophagus. The animals are then intubated with a cuffed endotracheal tube through the other nostril using a flexible fiberoptic bronchoscope as a guide. Pleural pressure is estimated with the esophageal balloon catheter (filled with one mL of air), which is positioned such that inspiration produces a negative pressure deflection with clearly discernible cardiogenic oscillations. Lateral pressure in the trachea is measured with a sidehole catheter (inner

dimension, 2.5 mm) advanced through and positioned distal to the tip of the nasotracheal tube. Transpulmonary pressure, the difference between tracheal pressure and pleural pressure, is measured with a differential pressure transducer (DP45; Validyne Corp., Northridge, CA). For the measurement of pulmonary resistance ( $R_L$ ), the maximal end of the nasotracheal tube is connected to a pneumotachograph (Fleisch, Dyna Sciences, Blue Bell, PA). The signals of flow and transpulmonary pressure are recorded on an oscilloscope (Model DR-12; Electronics for Medicine, White Plains, NY) which is linked to a PDP-11 Digital computer (Digital Equipment Corp., Maynard, MA) for on-line calculation of  $R_L$  from transpulmonary pressure, respiratory volume obtained by integration and flow. Analysis of 10-15 breaths is used for the determination of  $R_L$ . Thoracic gas volume ( $V_{tg}$ ) is measured in a body plethysmograph, to obtain specific pulmonary resistance ( $SR_L = R_L \cdot V_{tg}$ ).


The following examples are provided to illustrate the invention and are not to be construed as limiting the scope of the invention in any manner. In the examples, unless otherwise stated,

- all the end products of the formula I were analyzed by NMR, TLC and elementary analysis or mass spectroscopy;
- intermediates were analyzed by NMR and TLC;
- most compounds were purified by flash chromatography on silica gel, recrystallization and/or swish (suspension in a solvent followed by filtration of the solid);
- the course of reactions was followed by thin layer chromatography (TLC) and reaction times are given for illustration only;

the enantiomeric excess was measured on normal phase HPLC with a chiral column: ChiralPak AD; 250 x 4.6 mm.

#### EXAMPLE 1

[5-[(4-chlorophenyl)thio]-4-(methylsulfonyl)-6,7,8,9-tetrahydropyrido[3,2-b]indolizin-6-yl]acetic acid



**Step 1**      4-chloronicotinaldehyde

The title compound was prepared as described by F. Marsais et al., *J. Heterocyclic Chem.*, 25, 81 (1988).

**Step 2**      4-(methylthio)nicotinaldehyde

To a solution of NaSMe (9.5 g, 135 mmol) in MeOH (250 mL) was added the 4-chloronicotinaldehyde (13.5 g, 94.4 mmol) of Step 1 in MeOH (250 mL). The reaction mixture was maintained at 60°C for 15 min. The reaction mixture was poured over NH<sub>4</sub>Cl and EtOAc. The organic phase was separated, washed with H<sub>2</sub>O and dried over Na<sub>2</sub>SO<sub>4</sub>. The compound was then purified over silica gel with 50% EtOAc in Hexanes to provide the title compound.

**Step 3**      methyl (2Z)-2-azido-3-[4-(methylthio)pyridin-3-yl]prop-2-enoate

A solution of 4-(methylthio)nicotinaldehyde (4.8 g, 31 mmol) and methyl azidoacetate (9.0 g, 78 mmol) in MeOH (50 mL) was added to a solution of 25% NaOMe in MeOH (16.9 mL, 78 mmol) at -12°C. The internal temperature was monitored and maintained at -10°C to -12°C during the 30 min. addition. The resulting mixture was then stirred in an ice bath for several hours, followed by overnight in an ice bath in the cold room. The suspension was then poured onto a mixture of ice and NH<sub>4</sub>Cl, and the slurry was filtered after 10 min. of stirring. The product was washed with cold H<sub>2</sub>O and was then dried under vacuum to give the title compound as a beige solid (7.4 g), which contained some salts. The compound is then purified over silica gel with EtOAc.

**Step 4**      methyl 4-(methylthio)-1H-pyrrolo[2,3-b]pyridine-2-carboxylate

A suspension of the compound of Step 3 (0.40 g, 1.6 mmol) in xylenes (16 mL) was heated slowly to 140°C. After a period of 15 min. at 140°C, the yellow solution was cooled to room temperature. Precaution must be taken due to the possibility of an exotherme due to the formation of nitrogen. The suspension was then cooled to 0°C, filtered and washed with xylene to provide the title compound.

**Step 5**      ethyl 4-(methylthio)-6-oxo-6,7,8,9-tetrahydropyrido[3,2-b]indolizine-7-carboxylate

To a solution of the compound of Step 4 (0.35 g, 1.6 mmol) in DMF (20 mL) at 0°C was added NaH (1.2 eq.). After a period of 5 min., were added nBu<sub>4</sub>NI (0.10 g) and ethyl 4-bromobutyrate (0.40 mL). After a period of 1 h at room temperature, the reaction mixture was

poured over saturated  $\text{NH}_4\text{Cl}$  and EtOAc. The organic phase was separated, washed with  $\text{H}_2\text{O}$  and dried over  $\text{Na}_2\text{SO}_4$ . After evaporation the crude product was purified by flash chromatography. The bis ester was then dissolved in THF (7.0 mL) and a 1.06 M of THF solution of potassium tert-butoxide (2.2 mL) was added at  $0^\circ\text{C}$ . After a period of 1 h at room temperature, the reaction mixture was then poured over saturated  $\text{NH}_4\text{Cl}$  and EtOAc. The organic phase was separated, dried over  $\text{Na}_2\text{SO}_4$  and evaporated under reduced pressure to provide the title compound as a mixture of ethyl and methyl ester.

Step 6            4-(methylthio)-8,9-dihydropyrido[3,2-*b*]indolizin-6(7*H*)-one

To the compound of Step 5, (0.32 g) were added EtOH (8.0 mL) and concentrated HCl (2.0 mL). The resulting suspension was refluxed for 5 h. The reaction mixture was partitioned between EtOAc and  $\text{Na}_2\text{CO}_3$ . The organic phase was separated, evaporated to provide the title compound.

Step 7            ethyl (2*E*, 2*Z*)-[4-(methylthio)-8,9-dihydropyrido[3,2-*b*]indolizin-6(7*H*)-ylidene]ethanoate

To a DMF solution (12 mL) of triethyl phosphonoacetate (0.45 g, 2.17 mmol) were added 80% NaH (0.06 g, 2.00 mmol) and the compound of Step 6 (0.22 g, 1.00 mmole). After a period of 4 h at  $55^\circ\text{C}$ , the reaction mixture was poured over saturated  $\text{NH}_4\text{Cl}$  and EtOAc. The organic phase was separated and evaporated under reduced pressure. The crude product was purified by flash chromatography to afford the title compound.

Step 8            ethyl [4-(methylthio)-6,7,8,9-tetrahydropyrido[3,2-*b*]indolizin-6-yl]acetate

The compound of Step 7 was dissolved in MeOH – THF using heat for dissolution. To the previous cooled solution was added at room temperature  $\text{PtO}_2$  and the resulting mixture was maintained for 18 h under an atmospheric pressure of hydrogen. The reaction mixture was filtered carefully over celite using  $\text{CH}_2\text{Cl}_2$ . The filtrate was evaporated under reduced pressure to provide the title compound. Alternatively, the compounds of Step 7 can be hydrogenated with  $\text{Pd}(\text{OH})_2$  in EtOAc at 40 PSI of  $\text{H}_2$  for 18h.

Step 9            ethyl [4-(methylsulfonyl)-6,7,8,9-tetrahydropyrido[3,2-*b*]indolizin-6-yl]acetate

To the compound of Step 8 (0.08 g, 0.27 mmol) in MeOH (3.0 mL) were added  $\text{Na}_2\text{WO}_4$  (0.10 g) and 30%  $\text{H}_2\text{O}_2$  (600  $\mu\text{L}$ ). After a period of 1 h, the reaction mixture was

partitioned between H<sub>2</sub>O and EtOAc. The organic phase was washed with H<sub>2</sub>O, separated and evaporated. The title compound was purified by flash chromatography.

**Step 10**      ethyl [5-[(4-chlorophenyl)thio]-4-(methylsulfonyl)-6,7,8,9-tetrahydropyrido[3,2-*b*]indolizin-6-yl]acetate

To a 1,2-dichloroethane solution (2.0 mL) of 4,4'-dichlorodiphenyl disulfide (0.24 g) was added SO<sub>2</sub>Cl<sub>2</sub> (50  $\mu$ L). To the compound of Step 9 (0.05 g) in DMF (2.0 mL) was added the previous mixture ( $\approx$  180  $\mu$ L). The reaction was followed by <sup>1</sup>H NMR and maintained at room temperature until no starting material remained. The reaction mixture was poured over saturated NaHCO<sub>3</sub> and EtOAc. The organic phase was separated, evaporated and the title compound purified by flash chromatography.


**Step 11**      [5-[(4-chlorophenyl)thio]-4-(methylsulfonyl)-6,7,8,9-tetrahydropyrido[3,2-*b*]indolizin-6-yl]acetic acid

To the compound of Step 10 dissolved in a 1/1 mixture of THF-MeOH was added 1N NaOH. After a period of 18 h at room temperature, the reaction mixture was partitioned between saturated NH<sub>4</sub>Cl and EtOAc. The organic phase was separated, dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated to provide the title compound.

<sup>1</sup>H NMR (500 MHz, acetone-d<sub>6</sub>)  $\delta$  11.00 (bs, 1H), 8.60 (d, 1H), 7.80 (d, 1H), 7.20 (d, 2H), 7.00 (d, 2H), 4.65 (m, 1H), 4.20 (m, 1H), 3.75 (m, 1H), 3.35 (s, 3H), 2.80 to 2.10 (m, 6H).

**EXAMPLE 2**

[5-[(4-chlorophenyl)thio]-4-(methylthio)-6,7,8,9-tetrahydropyrido[3,2-*b*]indolizin-6-yl]acetic acid




The title compound can be prepared from the compound of Example 1, Step 8 in a similar manner as described in Example 1, Step 10 and 11.

m/z 418

EXAMPLE 3

[5-[(3,4-dichlorophenyl)thio]-4-(methylsulfonyl)-6,7,8,9-tetrahydropyrido[3,2-*b*]indolizin-6-yl]acetic acid (Enantiomer A and Enantiomer B)




The title compound was prepared as described in Example 1 using bis(3,4-dichlorophenyl)disulfide in Step 10.

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>) δ 8.55 (d, 1H), 7.85 (d, 1H), 7.35 (d, 1H), 7.15 (s, 1H), 6.95 (d, 1H), 4.60 (m, 1H), 4.15 (m, 1H), 3.80 (m, 1H), 3.40 (s, 3H), 2.80 to 2.10 (m, 6H).  
m/z 484.

The enantiomers were separated on a Chiralecel OD column 25 cm x 20 mm using 30 % isopropanol, 7 % ethanol, 0.2 % acetic acid in hexane, flow rate 8 ml/min. Their purities were verified on a Chiralecel OD column 25 cm x 4.6 mm using 35 % isopropanol 0.2 % acetic acid in hexane, flow rate 1.0 ml/min. More mobile enantiomer Tr = 9.7 min, less mobile enantiomer Tr 11.1 min.

EXAMPLE 4

[5-(4-chlorobenzoyl)-4-(methylsulfonyl)-6,7,8,9-tetrahydropyrido[3,2-*b*]indolizin-6-yl]acetic acid



**Step 1**      ethyl [5-(4-chlorobenzoyl)-4-(methylthio)-6,7,8,9-tetrahydropyrido[3,2-b]indolizin-6-yl]acetate

To a solution of 4-chlorobenzoyl chloride (0.30 g, 1.7 mmol) in 1,2-dichloroethane (6.0 mL) was added AlCl<sub>3</sub> (0.24 g, 1.8 mmole). After a period of 5 min. a solution of ethyl [4-(methylthio)-6,7,8,9-tetrahydropyrido[3,2-b]indolizin-6-yl]acetate from Example 1 Step 8 (0.15 g, 0.47 mmole) in 1,2-dichloroethane (6.0 mL) was added to the previous mixture. After a period of 4h, at 80°C, the reaction mixture was partitioned between EtOAc and NaHCO<sub>3</sub>. The organic phase was separated, dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated. The title compound was purified by flash chromatography.

**Step 2**      ethyl [5-(4-chlorobenzoyl)-4-(methylsulfonyl)-6,7,8,9-tetrahydropyrido [3,2-b]indolizin-6-yl]acetate

To a solution of ethyl[5-(4-chlorobenzoyl)-4-(methylthio)-6,7,8,9-tetrahydropyrido[3,2-b]indolizin-6yl] acetate (0.12 g, 0.27 mmole) in MeOH (5.0 mL) were added Na<sub>2</sub>WO<sub>4</sub> (0.1 g) and 30% H<sub>2</sub>O<sub>2</sub> (300 µL). The reaction mixture was stirred at 55°C for 1h. The reaction mixture was then partitioned between H<sub>2</sub>O and EtOAc. The organic phase was washed with H<sub>2</sub>O, dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated. The title compound was purified by flash chromatography.

**Step 3**      [5-(4-chlorobenzoyl)-4-(methylsulfonyl)-6,7,8,9-tetrahydropyrido[3,2-b]indolizin-6-yl]acetic acid


Ethyl [5-(4-chlorobenzoyl)-4-(methylsulfonyl)-6,7,8,9-tetrahydropyrido[3,2-b]indolizin-6yl]acetate was treated as described in Example 1 Step 11 to provide the title compound.

<sup>1</sup>H NMR (500 MHz, acetone-d<sub>6</sub>) δ 8.55 (d, 1H), 7.90 (d, 2H), 7.65 (d, 1H), 7.45 (d, 2H), 4.55 (m, 1H), 4.25 (m, 1H), 3.45 (m, 1H), 3.20 (s, 3H), 2.05 to 3.00 (m, 6H).  
m/z 446

**EXAMPLE 5**

[5-(4-bromophenyl)thio]-4-(methylsulfonyl)-6,7,8,9-tetrahydropyrido[3,2-b]indolizin-6-yl]acetic acid






The title compound was prepared as described in Example 1 using 4,4'-dibromodiphenyl disulfide.

<sup>1</sup>H NMR (500 MHz, Acetone-d<sub>6</sub>) δ 8.60 (d, 1H), 7.80 (d, 1H), 7.35 (d, 2H), 7.00 (d, 2H), 4.65 (m, 1H), 4.20 (m, 1H), 3.80 (m, 1H), 3.35 (s, 3H), 2.80 to 2.10 (m, 6H).

#### EXAMPLE 6 METHOD-1

[9-[(3,4-dichlorophenyl)thio]-1-(methylsulfonyl)-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-yl]acetic acid



#### Step 1      2-(methylthio)nicotinaldehyde

The title compound was prepared from 2-bromonicotinaldehyde (A. Numata *Synthesis* 1999 p.306) as described in Example 1 Step 2 except the solution was heated at 55°C for 2 hr.

#### Step 2      methyl (2Z)-2-azido-3-[2-(methylthio)pyridin-3-yl]prop-2-enoate

The title compound was prepared as described in Example 1 Step 3.

#### Step 3      methyl 4-(methylthio)-1H-pyrrolo[3,2-c]pyridine-2-carboxylate

A solution of methyl (2Z)-2-azido-3-[2-(methylthio)pyridin-3-yl]prop-2-enoate (1.00 g, 4.00 mmol) in mesitylene (50 mL) was heated at 160°C for a period of 1 h. The reaction mixture was cooled to room temperature then to 0°C, the precipitate was filtered and washed with cold mesitylene to provide the title compound.

**Step 4**      methyl 1-(methylthio)-8-oxo-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizine-7-carboxylate

To a suspension of methyl 4-(methylthio)-1H-pyrrolo[3,2-c]pyridine-2-carboxylate (0.30 g, 1.35 mmol) in THF (3 mL)- toluene (12.0 mL) were added a 1.06 M THF solution of potassium tert-butoxide (1.42 mL / 1.41 mmol) and methyl acrylate (300 µL). The resulting mixture was heated at 80°C for 18h. The mixture was partitioned between EtOAc and NH<sub>4</sub>Cl, filtered through celite. The organic phase was separated, dried over Na<sub>2</sub>SO<sub>4</sub> and filtered, to provide the title compound.

**Step 5**      1-(methylthio)-6,7-dihydro-8H-pyrido[3,4-b]pyrrolizin-8-one  
Methyl 1-(methylthio)-8-oxo-7,8-dihydro-6H-pyrido[3,4-b] pyrrolizine-7-carboxylate was converted to the title compound as described in Example 1 Step 6.

**Step 6**      methyl [8-hydroxy-1-(methylthio)-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-yl]acetate

A mixture of 1-(methylthio)-6,7-dihydro-8H-pyrido[3,4-b]pyrrolizin-8-one (0.15 g, 0.68 mmol), methyl bromoacetate (0.34 mL), Zn-Cu (0.226 g) in THF (3.0 mL) was sonicated for 2 h. The mixture was then heated at 60°C for 5 min. until completion of the reaction. The reaction mixture was partitioned between EtOAc and NH<sub>4</sub>Cl. The organic phase was separated, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and evaporated under reduced pressure to provide the title compound. The compound was purified by flash chromatography.

**Step 7**      methyl [1-(methylthio)-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-yl]acetate

To NaI (0.300 g) in CH<sub>3</sub>CN (3.2 mL) was added TMSCl (0.266 mL). This mixture was added to a suspension of methyl [8-hydroxy-1-(methylthio)-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-yl] acetate (0.15 g, 0.515 mmol) in CH<sub>3</sub>CN (1.5 mL), in a water bath. After a period of 0.5 h, the reaction mixture was partitioned between EtOAc and NaHCO<sub>3</sub>. The organic phase was separated, washed with sodium thiosulphate, dried over MgSO<sub>4</sub> and evaporated. The title compound was purified by flash chromatography.

**Step 8**      methyl [1-(methylsulfonyl)-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-yl]acetate

Methyl [1-(methylthio)-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-yl]acetate was converted to the title compound as described in Example 1 Step 9.

Step 9            [9-[(3,4-dichlorophenyl)thio]-1-(methylsulfonyl)-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-yl]acetic acid

Methyl [1-(methylsulfonyl)-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-yl]acetate was converted to the title compound as described in Example 1, Steps 10 and 11, using bis (3,4-dichlorophenyl)disulfide in Step 10.

<sup>1</sup>H NMR (500 MHz, acetone-d<sub>6</sub>) δ 8.35 (d, 1H) 7.80 (d, 1H), 7.35 (d, 1H), 7.15 (s, 1H), 6.95 (d, 1H), 4.55 (m, 1H), 4.35 (m, 1H), 3.90 (m, 1H), 3.30 (s, 3H), 3.15 (m, 1H), 3.05 (m, 1H), 2.80 (m, 1H), 2.50 (m, 1H).

EXAMPLE 6 METHOD-2

[9-[(3,4-dichlorophenyl)thio]-1-(methylsulfonyl)-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-yl]acetic acid

Step 1            1-(methylthio)-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-ol

To a suspension of 1-(methylthio)-6,7-dihydro-8H-pyrido[3,4-b]pyrrolizin-8-one from Example 6 method-1 Step 5 (0.55 g, 2.2 mmol) in EtOH (10 mL)-THF (1 mL) was added NaBH<sub>4</sub> (0.10 g, 2.6 mmol) at 0°C. After a period of 30 min. at room temperature, the reaction was quenched by the addition of acetone. The solvents were evaporated under reduced pressure and EtOAc and H<sub>2</sub>O were added to the residue. The organic phase was separated, dried over MgSO<sub>4</sub> and evaporated. The title compound was washed with EtOAc/Hexane and filtered.

Step 2            dimethyl 2-[1-(methylthio)-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-yl]malonate

To a suspension of 1-(methylthio)-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-ol (0.54 g, 2.1 mmol) in THF (10 mL) at -78°C were added 1M NaHMDS in THF (2.35 mL, 2.4 mmol) and diphenyl chlorophosphate (0.53 mL, 2.6 mmol). After a period of 30 min. dimethyl malonate (0.73 mL, 6.4 mmol) and 1M NaHMDS in THF (6.8 mL, 6.8 mmol) were added. The reaction mixture was brought to 0°C and then to room temperature. The mixture was then partitioned between EtOAc and NH<sub>4</sub>Cl. The organic phase was dried over MgSO<sub>4</sub>, filtered and evaporated. The title compound was purified by flash chromatography.

Step 3            methyl [1-(methylthio)-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-yl]-acetate

To a mixture of dimethyl 2-[1-(methylthio)-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-yl]malonate (0.59 g, 2.17 mmol) and DMSO (4mL) was added NaCl (0.45 g) in H<sub>2</sub>O (0.45 mL). After a period of 18 h at 150°C, the reaction mixture was partitioned between


ETOAc and H<sub>2</sub>O. The organic phase was separated, dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated. The title compound was then purified by flash chromatography.

Step 4      [9-[(3,4-dichlorophenyl)thio]-1-(methylsulfonyl)-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-yl]acetic acid

The title compound was obtained from methyl [1-(methylthio)-7,8-dihydro-6H-pyrido-[3,4-b]pyrrolizin-8yl]acetate as described in Example 6 Method-1 Step 8 to 9.

EXAMPLE 7

[10-[(3,4-dichlorophenyl)sulfanyl]-1-(methylsulfonyl)-6,7,8,9-tetrahydropyrido[3,4-b]indolizin-9-yl]acetic acid



Step 1      ethyl [1-(methylsulfonyl)-6,7,8,9-tetrahydropyrido[3,4-b]indolizin-9-yl]acetate

The title compound was prepared from the product of Example 6 Step 3 in the same manner as described in Example 1 Steps 5 to 9.


Step 2      [10-[(3,4-dichlorophenyl)sulfanyl]-1-(methylsulfonyl)-6,7,8,9-tetrahydropyrido[3,4-b]indolizin-9-yl]acetic acid

The product of Step 1 was converted to the title compound in the same manner as Example 1, Steps 10-11, using bis (3,4-dichlorophenyl)disulfide in Step 10.

MS M+1=485

EXAMPLE 8

(4-(methylsulfonyl)-5-{[4-(trifluoromethyl)phenyl]thio}-6,7,8,9-tetrahydropyrido[3,2-b]indolizin-6-yl)acetic acid




The title compound was prepared as described in Example 1 using bis[4-trifluoromethylphenyl]disulfide

<sup>1</sup>H NMR (500 MHz, acetone-d<sub>6</sub>) δ 8.55 (d, 1H), 7.75 (d, 1H), 7.45 (d, 2H), 7.15 (d, 2H), 4.55 (m, 1H), 4.15 (m, 1H), 3.80 (m, 1H), 3.30 (s, 3H), 2.80 to 2.10 (m, 6H).

m/z 513 (M+1)

#### EXAMPLE 9

[5-[(2-chloro-4-fluorophenyl)thio]-4-(methylsulfonyl)-6,7,8,9-tetrahydropyrido[3,2-b]indolizin-6-yl]acetic acid




The title compound was prepared as described in Example 1 using bis(2-chloro-4-fluorophenyl)disulfide.

m/z 469 (M+1)

#### EXAMPLE 10

[4-(methylsulfonyl)-5-(2-naphthylthio)-6,7,8,9-tetrahydropyrido[3,2-b]indolizin-6-yl]acetic acid




The title compound was prepared as described in Example 1 using di(2-naphthyl) disulfide.

M/z 467 (M+1)

#### EXAMPLE 11

[5-[(2,3-dichlorophenyl)thio]-4-(methylsulfonyl)-6,7,8,9-tetrahydropyrido[3,2-b]indolizin-6-yl]acetic acid




The title compound was prepared as described in Example 1 using bis(2,3-dichlorophenyl)disulfide.

$^1\text{H}$  NMR (500 MHz, acetone- $d_6$ )  $\delta$  8.85 (d, 1H), 7.80 (d, 1H), 7.30 (d, 1H), 7.00 (t, 1H), 6.60 (d, 1H), 4.60 (m, 1H), 4.20 (m, 1H), 3.80 (m, 1H), 3.40 (s, 3H), 2.80 to 2.10 (m, 6H).

#### EXAMPLE 12

[5-[(4-methylphenyl)thio]-4-(methylsulfonyl)-6,7,8,9-tetrahydropyrido[3,2-b]indolizin-6-yl]acetic acid




The title compound was prepared as described in Example 1 using p-tolyl disulfide.

$^1\text{H}$  NMR (500 MHz, acetone- $d_6$ )  $\delta$  8.55 (d, 1H), 7.80 (d, 1H), 6.95 (m, 4H), 4.60 (m, 1H), 4.15 (m, 1H), 3.80 (m, 1H), 3.35 (s, 3H), 2.80 to 2.10 (m, 6H).

#### EXAMPLE 13

[4-(methylsulfonyl)-5-(phenylthio)-6,7,8,9-tetrahydropyrido[3,2-*b*]indolizin-6-yl]acetic acid




The title compound was prepared as described in Example 1 using diphenyl disulfide.

$^1\text{H}$  NMR (500 MHz, acetone- $d_6$ )  $\delta$  8.55 (d, 1H), 7.80 (d, 1H), 7.15 to 6.90 (m, 5H), 4.60 (m, 1H), 4.15 (m, 1H), 3.75 (m, 1H), 3.30 (s, 3H), 2.80 to 2.10 (m, 6H).

#### EXAMPLE 14

[5-[(2,4-dichlorophenyl)thio]-4-(methylsulfonyl)-6,7,8,9-tetrahydropyrido[3,2-*b*]indolizin-6-yl]acetic acid




The title compound was prepared as described in Example 1 using bis(2,4-dichlorophenyl)disulfide. The disulfide was prepared from 2,4-dichlorothiophenyl using Br<sub>2</sub> in ether.

<sup>1</sup>H NMR (500 MHz, acetone-d<sub>6</sub>) δ 8.55 (d, 1H), 7.85 (d, 1H), 7.35 (s, 1H), 7.00 (d, 1H), 6.65 (d, 1H), 4.55 (m, 1H), 4.15 (m, 1H), 3.80 (m, 1H), 3.35 (s, 3H), 2.80 to 2.10 (m, 6H).

#### EXAMPLE 15

[5-[(4-chlorophenyl)thio]-4-(methylsulfonyl)-6,7,8,9-tetrahydropyrido[4,3-*b*]indolizin-6-yl]acetic acid




The title compound was prepared as described in Example 1 from 3-chloronicotinaldehyde (Heterocycles p. 151, 1993) except the thermal cyclization was performed by adding the azide to decalin at reflux.

<sup>1</sup>H NMR (500 MHz, acetone-d<sub>6</sub>) δ 9.20 (s, 1H), 8.85 (s, 1H), 7.20 (d, 2H), 7.00 (d, 2H), 4.70 (m, 1H), 4.30 (m, 1H), 3.75 (m, 1H), 3.35 (s, 3H), 2.80 to 2.10 (m, 6H).



**EXAMPLE 16**

[9-[(4-chlorophenyl)thio]-1-(methylsulfonyl)-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-yl]acetic acid




The title compound was prepared from the product of Example 6 Method 1 Step 8, as described in the procedures outlined in Example 1 Steps 10 and 11, using bis (4-chlorophenyl)disulfide in Step 10.

<sup>1</sup>H NMR (500 MHz, acetone-d<sub>6</sub>) δ 8.25-8.3 (m, 1H), 7.71-7.75 (m, 1H), 7.12-7.17 (m, 2H), 6.97-7.04 (m, 2H), 4.45-4.51 (m, 1H), 4.32-4.39 (m, 1H), 3.73-3.80 (m, 1H), 3.29 (s, 3H), 3.15-3.21 (m, 1H), 2.99-3.08 (m, 1H), 2.66-2.73 (m, 1H), 2.46-2.54 (m, 1H).

**EXAMPLE 17**

{9-[(3,4-dichlorophenyl)thio]-1-isopropyl-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-yl}acetic acid (Enantiomer A and Enantiomer B)

**Step 1** 2-chloronicotinaldehyde

To a solution of diisopropyl amine (110 mL, 780 mmol) in THF (500 mL) was added a 2.5 M hexanes solution of n-BuLi (300 mL, 750 mmol) at -40°C. After 5 min, the reaction mixture was cooled to -95°C then DMPU (15 mL) and 2-chloropyridine (50 mL, 532 mmol) were successively added. The resulting mixture was then warmed and stirred at -78°C for 4h. After this time, the yellow suspension was cooled again to -95°C before DMF (70 mL)

was added. The final reaction mixture was warmed to  $-78^{\circ}\text{C}$  and stirred at that temperature for 1.5h. The reaction mixture was poured into cold aqueous HCl (3N, 800 mL) and stirred for 5 min. Aqueous concentrated  $\text{NH}_4\text{OH}$  was added to adjust pH to 7.5. The aqueous layer was extracted three times with EtOAc. The combined organic layer was washed with aqueous  $\text{NH}_4\text{Cl}$  and brine, dried over anhydrous  $\text{Na}_2\text{SO}_4$ , filtered and concentrated. The crude material was further purified by a pad of silica gel by eluting with a gradient from 100% hexanes to 100% EtOAc and the product was crystallized in cold hexanes to yield the title compound as a pale yellow solid.

**Step 2**            methyl (2Z)-2-azido-3-(2-chloropyridin-3-yl)prop-2-enoate

The title compound was prepared as described in Example 1 Step 3.

**Step 3**            methyl 4-chloro-1H-pyrrolo[3,2-c]pyridine-2-carboxylate

The title compound was prepared in a similar manner as described in Example 6 Method-1 Step 3.

**Step 4**            methyl 1-chloro-8-oxo-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizine-7-carboxylate

To a suspension of methyl 4-chloro-1H-pyrrolo[3,2-c]pyridine-2-carboxylate (12.5 g, 59 mmol) in THF (116 mL) – toluene (460 mL) were added a 1.0 M THF solution of potassium tert-butoxide (64 mL, 64 mmol) and methyl acrylate (55 mL, 611 mmol). The resulting mixture was heated at  $100^{\circ}\text{C}$  for 18h. After this time, the suspension was cooled to room temperature and it was poured into a mixture of saturated aqueous  $\text{NH}_4\text{Cl}$  (400 mL) and hexanes (400 mL). The solids were decanted, filtered and washed with  $\text{H}_2\text{O}$  and hexanes to provide the title compound.

**Step 5**            1-chloro-6,7-dihydro-8H-pyrido[3,4-b]pyrrolizin-8-one

The title compound was prepared in a similar manner as described in Example 1 Step 6 using isopropanol instead of ethanol and heating at  $100^{\circ}\text{C}$  for 1h.

**Step 6**            1-isopropenyl-6,7-dihydro-8H-pyrido[3,4-b]pyrrolizin-8-one

To a mixture of 1-chloro-6,7-dihydro-8H-pyrido[3,4-b]pyrrolizin-8-one (5.0 g, 24.3 mmol), tris (dibenzylidene acetone)dipalladium (0) (1.0 g, 1.09 mmol) and triphenylarsine (2.70 g, 8.82 mmol) in DMF (100 mL) was added tributylisopropenyl stannane (9.60 g, 29.00 mmol). The resulting mixture was degassed and heated at  $78^{\circ}\text{C}$  for a period of 18 h. The

solvent was evaporated under reduced pressure.  $\text{CH}_2\text{Cl}_2$  and celite were added to the resulting mixture which was then filtered over celite. The title compound was purified by flash chromatography (50% to 100% EtOAc in Hexane).

Step 7            ethyl (2E)-(1-isopropenyl-6,7-dihydro-8H-pyrido[3,4-b]pyrrolizin-8-ylidene)ethanoate

To a solution of 1-isopropenyl-6,7-dihydro-8H-pyrido[3,4-b]pyrrolizin-8-one (0.60 g, 2.8 mmol) and triethyl phosphonoacetate (1.00 g, 4.46 mmol) in THF (24 mL) at  $-78^\circ\text{C}$  was added 80% NaH (0.12 g, 4.00 mmol), the reaction mixture was allowed to warm to  $0^\circ\text{C}$ , then to room temperature. The reaction mixture was poured onto saturated  $\text{NH}_4\text{Cl}$  and EtOAc. The organic phase was separated, dried over  $\text{Na}_2\text{SO}_4$  and evaporated. The title compound was purified by flash chromatography (40% EtOAc in Hexane).

Step 8            ethyl (1-isopropyl-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-yl)acetate

To a solution of ethyl (2E)-(1-isopropenyl-6,7-dihydro-8H-pyrido[3,4-b]pyrrolizin-8-ylidene)ethanoate (0.40 g, 1.4 mmol) in MeOH (20 mL) was added  $\text{Pd}(\text{OH})_2$  (0.20 g). The mixture was stirred under 1 atm of  $\text{H}_2$  for 3h. The mixture was filtered over celite and evaporated to provide the title compound.

Step 9            ethyl {9-[(3,4-dichlorophenyl)thio]-1-isopropyl-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-yl}acetate

To a solution of bis (3,4-dichlorophenyl)disulfide (0.24 g, 0.67 mmol) in  $\text{CH}_2\text{Cl}_2$  (5.6 mL) was added  $\text{SO}_2\text{Cl}_2$  (0.036 mL). The resulting yellow mixture was stirred at room temperature for 1 h. This solution was added to a solution of ethyl (1-isopropyl-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-yl) acetate (0.15 g, 0.52 mmol) in DMF (5.6 mL) at  $0^\circ\text{C}$ . After 1.5 h at  $0^\circ\text{C}$ , the reaction mixture was poured over saturated  $\text{NaHCO}_3$  and EtOAc. The organic phase was separated, dried over  $\text{Na}_2\text{SO}_4$ , filtered and evaporated. The title compound was purified by flash chromatography (30% to 40% EtOAc in Hexane).

Step 10            {9-[(3,4-dichlorophenyl)thio]-1-isopropyl-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-yl}acetic acid

To a solution of ethyl {9-[(3,4-dichlorophenyl)thio]-1-isopropyl-7,8-dihydro-6H-pyrido[3,4-b]pyrrolizin-8-yl}acetate (0.23 g, 0.50 mmol) in THF (5 mL) and MeOH (2.5 mL) was added 1.0 M NaOH (1.5 mL, 1.5 mmol). After stirring 18h at RT, HOAc (0.25 mL) was added and the solvent was evaporated. The residue was taken up in EtOAc/ $\text{H}_2\text{O}$ , and the organic layer


was washed with H<sub>2</sub>O and brine. After drying (Na<sub>2</sub>SO<sub>4</sub>), the solution was filtered and evaporated. The residue was stirred with 1:1 EtOAc:hex to give, after filtration, the title compound as a white solid.

<sup>1</sup>H NMR (MeOH-d<sub>4</sub>) δ 1.14–1.26 (m, 6H), 2.47–2.56 (m, 1H), 2.56–2.64 (m, 1H), 2.94–3.05 (m, 2H), 3.81–3.89 (m, 1H), 4.22–4.30 (m, 1H), 4.33–4.44 (m, 2H), 6.93–6.99 (m, 1H), 7.14–7.19 (m, 1H), 7.33–7.39 (m, 1H), 7.54–7.59 (m, 1H), 8.16–8.21 (m, 1H).

The product of Step 10 was converted to its methyl ester using CH<sub>2</sub>N<sub>2</sub>, and the ester was subjected to HPLC separation on chiral stationary phase (chiralcel OD column 2x25cm), eluting with 12% 2-propanol in hexane at a flow rate of 6 mL/min. Enantiomer A (less polar) has a retention time of 31.9 min and Enantiomer B (more polar) has a retention time of 35.5 min. Both A and B were hydrolyzed as in Ex. 17 Step 10 to give enantiomers A and B of the title compound.

## WHAT IS CLAIMED IS:

1. A compound having the formula I



I

and pharmaceutically acceptable salts and hydrates thereof, wherein:

A is selected from C<sub>1-3</sub>alkyl optionally substituted with one to four halogen atoms, O(CH<sub>2</sub>)<sub>1-2</sub>, and S(CH<sub>2</sub>)<sub>1-2</sub>;

Ar is aryl or heteroaryl each optionally substituted with one to four groups independently selected from R<sub>G</sub>;

Q is selected from:

- (1) COOH,
- (2) CONR<sup>a</sup>R<sup>b</sup>,
- (3) C(O)NHSO<sub>2</sub>R<sup>c</sup>,
- (4) SO<sub>2</sub>NHR<sup>a</sup>,
- (5) SO<sub>3</sub>H,
- (6) PO<sub>3</sub>H<sub>2</sub>, and
- (7) tetrazolyl;

one of X<sup>1</sup>, X<sup>2</sup>, X<sup>3</sup> or X<sup>4</sup> is nitrogen and the others are independently selected from CH and C-R<sub>G</sub>;

Y<sup>1</sup> is selected from -(CR<sup>d</sup>Re)<sub>a</sub>-X-(CR<sup>d</sup>Re)<sub>b</sub>-, phenylene, C<sub>3-6</sub>cycloalkylidene and

C<sub>3-6</sub>cycloalkylene, wherein a and b are integers 0-1 such that the sum of a and b equals 0, 1 or 2,

and X is a bond, O, S, NR<sup>a</sup>, C(O), CH(OR<sup>a</sup>), OC(O), C(O)O, C(O)NR<sup>a</sup>, OC(O)NR<sup>a</sup>, NR<sup>a</sup>C(O), CR<sup>d</sup>=CR<sup>e</sup> or C≡C;

Y<sup>2</sup> is selected from (CR<sup>d</sup>Re)<sub>m</sub> and CR<sup>d</sup>=CR<sup>e</sup>;

R<sup>1</sup> is selected from H, CN, OR<sup>a</sup>, S(O)<sub>n</sub>C<sub>1-6</sub>alkyl and C<sub>1-6</sub>alkyl optionally substituted with one to six groups independently selected from halogen, OR<sup>a</sup> and S(O)<sub>n</sub>C<sub>1-6</sub>alkyl;

R<sup>2</sup> is selected from H and C<sub>1-6</sub>alkyl optionally substituted with one to six halogen; or

R<sup>1</sup> and R<sup>2</sup> together represent an oxo; or

R<sup>1</sup> and R<sup>2</sup> taken together form a 3- or 4- membered ring containing 0 or 1 heteroatom selected from NR<sup>f</sup>, S, and O optionally substituted with one or two groups selected from F, CF<sub>3</sub> and CH<sub>3</sub>;

R<sup>3</sup> is selected from H and C<sub>1-6</sub>alkyl optionally substituted with one to six groups independently selected from OR<sup>a</sup> and halogen;

R<sup>a</sup> and R<sup>b</sup> are independently selected from H, C<sub>1-10</sub>alkyl, C<sub>2-10</sub>alkenyl, C<sub>2-10</sub>alkynyl, Cy and Cy C<sub>1-10</sub>alkyl, wherein said alkyl, alkenyl, alkynyl and Cy are optionally substituted with one to six substituents independently selected from halogen, amino, carboxy, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, aryl, heteroaryl, aryl C<sub>1-4</sub>alkyl, hydroxy, CF<sub>3</sub>, OC(O)C<sub>1-4</sub>alkyl, OC(O)NR<sup>i</sup>R<sup>j</sup>, and aryloxy; or R<sup>a</sup> and R<sup>b</sup> together with the atom(s) to which they are attached form a heterocyclic ring of 4 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N-R<sup>f</sup>;

R<sup>c</sup> is selected from C<sub>1-6</sub>alkyl optionally substituted with one to six halogen, aryl and heteroaryl, wherein said aryl and heteroaryl are optionally substituted with one to three groups selected from halogen, OC<sub>1-6</sub>alkyl, O-haloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl and haloC<sub>1-6</sub>alkyl;

R<sup>d</sup> and R<sup>e</sup> are independently H, halogen, aryl, heteroaryl, C<sub>1-6</sub>alkyl or haloC<sub>1-6</sub>alkyl;

R<sup>f</sup> is selected from H, C<sub>1-6</sub>alkyl, haloC<sub>1-6</sub>alkyl, Cy, C(O)C<sub>1-6</sub>alkyl, C(O)haloC<sub>1-6</sub>alkyl, and C(O)-Cy;

R<sup>g</sup> is selected from

- (1) halogen,
- (2) CN,
- (3) C<sub>1-6</sub>alkyl optionally substituted with one to eight groups independently selected from aryl, heteroaryl, halogen, NR<sup>a</sup>R<sup>b</sup>, C(O)R<sup>a</sup>, C(OR<sup>a</sup>)R<sup>a</sup>R<sup>b</sup>, SR<sup>a</sup> and OR<sup>a</sup>, wherein aryl, heteroaryl and alkyl are each optionally substituted with one to six groups independently selected from halogen, CF<sub>3</sub>, and COOH,
- (4) C<sub>2-6</sub>alkenyl optionally substituted with one to six groups independently selected from halogen and OR<sup>a</sup>,
- (5) Cy
- (6) C(O)R<sup>a</sup>,

- (7)  $C(O)OR^a$ ,
- (8)  $CONR^aR^b$ ,
- (9)  $OC(=O)NR^aR^b$ ,
- (10)  $OC_{1-6}alkyl$ , wherein alkyl is optionally substituted with one to six substituents selected from halogen, aryl, heteroaryl, OH and  $OC(O)R^a$ ,
- (11) O-Cy,
- (12)  $S(O)_nC_{1-6}alkyl$ , wherein alkyl is optionally substituted with one to six substituents selected from halogen, aryl, heteroaryl, OH, and  $OC(O)R^a$ ,
- (13)  $S(O)_n-Cy$ ,
- (14)  $-NR^aS(O)_nR^b$ ,
- (15)  $-NR^aR^b$ ,
- (16)  $-NR^aC(O)R^b$ ,
- (17)  $-NR^aC(O)OR^b$ ,
- (18)  $-NR^aC(O)NR^aR^b$ ,
- (19)  $S(O)_nNR^aR^b$ ,
- (20)  $NO_2$ ,
- (21)  $C_{5-8}cycloalkenyl$ ,

wherein Cy is optionally substituted with one to eight groups independently selected from halogen,  $C(O)R^a$ ,  $OR^a$ ,  $C_{1-3}alkyl$ , aryl, heteroaryl and  $CF_3$ ;

$R^i$  and  $R^j$  are independently selected from hydrogen,  $C_{1-10}alkyl$ , Cy and  $Cy-C_{1-10}alkyl$ ; or

$R^i$  and  $R^j$  together with the nitrogen atom to which they are attached form a ring of 5 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and N- $R^f$ ;


Cy is selected from heterocyclyl, aryl, and heteroaryl;

m is 1, 2 or 3; and

n is 0, 1 or 2.

- 2. A compound of Claim 1 wherein A-Q is  $CH_2CO_2H$ .
- 3. A compound of Claim 1 wherein Ar is naphthyl or optionally substituted phenyl wherein said substituents are 1 or 2 groups independently selected from  $R^g$ .
- 4. A compound of Claim 1 wherein  $Y^1$  is selected from  $C(O)$  and S.

5. A compound of Claim 1 wherein one of  $X^1$ ,  $X^2$  and  $X^3$  is nitrogen and the others are independently CH or CR<sub>g</sub>, and  $X^4$  is CR<sub>g</sub>.
6. A compound of Claim 1 wherein one of  $X^1$ ,  $X^2$  and  $X^3$  is nitrogen and the others are CH, and  $X^4$  is C-S(O)<sub>n</sub>-C<sub>1-6</sub>alkyl or C-C<sub>1-6</sub>alkyl optionally substituted with OR<sup>a</sup>.
7. A compound of Claim 1 wherein R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> are each hydrogen.
8. A compound of Claim 1 wherein Y<sup>2</sup> is selected from CH<sub>2</sub> and CH<sub>2</sub>CH<sub>2</sub>.
9. A compound of Claim 1 represented by the formula Ia:




Ia

wherein  $X^2$  and  $X^3$  are independently CH or C-R<sub>g</sub>, A, Ar, Q, Y<sup>1</sup>, R<sup>1</sup>, R<sup>2</sup>, m and R<sub>g</sub> are as defined in Claim 1.

10. A compound of Claim 9 wherein  $X^2$  and  $X^3$  are each CH, R<sup>1</sup> and R<sup>2</sup> are each H, and A-Q is CH<sub>2</sub>CO<sub>2</sub>H.
11. A compound of Claim 9 wherein Y<sup>1</sup>-Ar is S-phenyl optionally substituted with 1 or 2 groups independently selected from halogen, C<sub>1-6</sub> alkyl and trifluoromethyl.
12. A compound of Claim 1 represented by the formula Ib:






wherein  $X^1$  and  $X^2$  are independently CH or C-Rg, A, Ar, Q,  $Y^1$ ,  $R^1$ ,  $R^2$ , m and Rg are as defined in Claim 1.

13. A compound of Claim 12 wherein  $X^1$  and  $X^2$  are each CH,  $R^1$  and  $R^2$  are each H, and A-Q is  $CH_2CO_2H$ .

14. A compound of Claim 13 wherein  $Y^1$ -Ar is S-phenyl optionally substituted with 1 or 2 groups independently selected from halogen,  $C_{1-6}$  alkyl and trifluoromethyl.


15. A compound of Claim 1 represented by the formula Ic:



wherein one of  $X^1$ ,  $X^2$  and  $X^3$  is N and the others are each CH,  $X^4$  is CRg, m is 1 or 2, and Ar,  $Y^1$  and m are as defined in Claim 1.

16. A compound of Claim 15 wherein Ar is phenyl optionally substituted with 1 or 2 groups independently selected from halogen,  $C_{1-3}$ alkyl and trifluoromethyl.


17. A compound of Claim 15 wherein  $Y^1$  is S or C(O).
18. A compound of Claim 15 wherein  $X^4$  is selected from C-S(O)<sub>n</sub>-C<sub>1-6</sub>alkyl and C-C<sub>1-6</sub>alkyl optionally substituted with OR<sup>a</sup>.
19. A compound of Claim 15 wherein  $Y^1$ -Ar is S-phenyl optionally substituted with 1 or 2 groups independently selected from halogen, C<sub>1-6</sub>alkyl and trifluoromethyl;  $X^1$  and  $X^2$  are each CH,  $X^3$  is N, m is 1 or 2, and  $X^4$  is C-SO<sub>2</sub>C<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyl.
20. A compound of Claim 1 selected from:



$X^1$	$X^2$	$X^3$	$X^4$	Ar	$Y^1$	m
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	4-Cl-Ph	S	2
N	CH	CH	C(SCH <sub>3</sub> )	4-Cl-Ph	S	2
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	3,4-diCl-Ph	S	2
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	4-Cl-Ph	C(O)	2
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	4-Br-Ph	S	2
CH	CH	N	C(SO <sub>2</sub> CH <sub>3</sub> )	3,4-diCl-Ph	S	1
CH	CH	N	C(SO <sub>2</sub> CH <sub>3</sub> )	3,4-diCl-Ph	S	2
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	4-CF <sub>3</sub> -Ph	S	2
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	2-Cl-4-F-Ph	S	2
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	2-naphthyl	S	2
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	2,3-diCl-Ph	S	2
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	4-CH <sub>3</sub> -Ph	S	2
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	Ph	S	2
N	CH	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	2,4-diCl-Ph	S	2
CH	N	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	4-Cl-Ph	S	2

X1	X2	X3	X4	Ar	Y1	m
CH	CH	N	C(SO <sub>2</sub> CH <sub>3</sub> )	4-Cl-Ph	S	2
N	C(CH <sub>3</sub> )	CH	C(SO <sub>2</sub> CH <sub>3</sub> )	4-Cl-Ph	S	2
N	CH	C(CH <sub>3</sub> )	C(SO <sub>2</sub> CH <sub>3</sub> )	4-Cl-Ph	S	2
CH	C(CH <sub>3</sub> )	N	C(SO <sub>2</sub> CH <sub>3</sub> )	4-Cl-Ph	S	2
C(CH <sub>3</sub> )	CH	N	C(SO <sub>2</sub> CH <sub>3</sub> )	4-Cl-Ph	S	2
N	CH	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-F-Ph	S	2
N	CH	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-Cl-Ph	S	2
N	CH	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	2,4-diCl-Ph	S	2
N	CH	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-Br-Ph	S	2
N	CH	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	2-Cl-4-F-Ph	S	2
N	CH	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	3,4-diCl-Ph	S	2
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-F-Ph	S	2
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-Cl-Ph	S	2
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	2,4-diCl-Ph	S	2
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-Br-Ph	S	2
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	2-Cl-4-F-Ph	S	2
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	3,4-diCl-Ph	S	2
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-F-Ph	S	1
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-Cl-Ph	S	1
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	2,4-diCl-Ph	S	1
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-Br-Ph	S	1
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	2-Cl-4-F-Ph	S	1
CH	CH	N	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	3,4-diCl-Ph	S	1
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-F-Ph	S	1
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-Cl-Ph	S	1
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	2,4-diCl-Ph	S	1
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-Br-Ph	S	1
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	2-Cl-4-F-Ph	S	1
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	3,4-diCl-Ph	S	1
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-F-Ph	S	2
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-Cl-Ph	S	2
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	2,4-diCl-Ph	S	2

X <sup>1</sup>	X <sup>2</sup>	X <sup>3</sup>	X <sup>4</sup>	Ar	Y <sup>1</sup>	m
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	4-Br-Ph	S	2
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	2-Cl-4-F-Ph	S	2
CH	N	CH	C(CH(CH <sub>3</sub> ) <sub>2</sub> )	3,4-diCl-Ph	S	2
N	CH	CH	C(CH(OCH <sub>3</sub> ) (CH <sub>2</sub> CH <sub>3</sub> ))	4-Cl-Ph	S	2
N	CH	CH	C(CH(OCH <sub>3</sub> ) (CH <sub>2</sub> CH <sub>3</sub> ))	4-Cl-Ph	S	1
CH	N	CH	C(CH(OCH <sub>3</sub> ) (CH <sub>2</sub> CH <sub>3</sub> ))	4-Cl-Ph	S	1
CH	N	CH	C(CH(OCH <sub>3</sub> ) (CH <sub>2</sub> CH <sub>3</sub> ))	4-Cl-Ph	S	2
CH	CH	N	C(CH(OCH <sub>3</sub> ) (CH <sub>2</sub> CH <sub>3</sub> ))	4-Cl-Ph	S	2
CH	CH	N	C(CH(OCH <sub>3</sub> ) (CH <sub>2</sub> CH <sub>3</sub> ))	4-Cl-Ph	S	1
N	CH	CH	C(C(CH <sub>3</sub> ) <sub>3</sub> )	4-Cl-Ph	S	2
N	CH	CH	C(C(CH <sub>3</sub> ) <sub>3</sub> )	3,4-diCl-Ph	S	2
N	CH	CH	C(C(CH <sub>3</sub> ) <sub>3</sub> )	4-Br-Ph	S	2
N	CH	CH	C(C(CH <sub>3</sub> ) <sub>3</sub> )	4-CF <sub>3</sub> -Ph	S	2
N	CH	CH	C(C(CH <sub>3</sub> ) <sub>3</sub> )	2-Cl-4-F-Ph	S	2
N	CH	CH	C(C(CH <sub>3</sub> ) <sub>3</sub> )	2-naphthyl	S	2
N	CH	CH	C(C(CH <sub>3</sub> ) <sub>3</sub> )	2,3-diCl-Ph	S	2
N	CH	CH	C(C(CH <sub>3</sub> ) <sub>3</sub> )	4-CH <sub>3</sub> -Ph	S	2
N	CH	CH	C(C(CH <sub>3</sub> ) <sub>3</sub> )	Ph	S	2
N	CH	CH	C(C(CH <sub>3</sub> ) <sub>3</sub> )	2,4-diCl-Ph	S	2



Ar	Y <sup>1</sup>
----	----------------

Ar	Y1
5-tetrazolyl	S
2-pyrrolyl	S
1,2,4-triazoly-3-yl	S
1,2,3-triazol-4-yl	S
5-imidazolyl	S
4-pyrazolyl	S
5-pyrazolyl	S
(1H,4H)-5-oxo-1,2,4-triazol-3-yl	S
4-isothiazolyl	S
1,2,5-thiadiazol-5-yl	S
1,2,5-oxadiazol-5-yl	S
3-furanyl	S
1,2,3-thiadiazol-4-yl	S
1,2,3-oxadiazol-4-yl	S
4-isoxazolyl	S
3-thienyl	S
4-oxazolyl	S
4-thiazolyl	S
(5H)-2-oxo-5-furanyl	S
(5H)-2-oxo-4-furanyl	S
1,2,4-oxadiazol-5-yl	S
3-pyridyl	S
2-pyrazinyl	S
5-pyrimidinyl	S
2-indolyl	S
2-benzothienyl	S
2-benzofuranyl	S
4-oxo-benzopyran-2-yl	S
2-quinolinyl	S
2-benzimidazolyl	S
2-benzoxazolyl	S
2-benzothiazolyl	S

Ar	Y1
1-benzotriazolyl	CH <sub>2</sub> S
thieno[2,3-b]pyridin-2-yl	S

21. A pharmaceutical composition comprising a compound of formula I as defined in any one of Claims 1 to 20, or a pharmaceutically acceptable salt or hydrate thereof, and a pharmaceutically acceptable carrier.

22. The composition of Claim 21 further comprising a second active ingredient selected from an antihistamine, a leukotriene antagonist and a leukotriene biosynthesis inhibitor.

23. A method for the treatment of prostaglandin D<sub>2</sub> mediated diseases which comprises administering to a patient in need of such treatment a therapeutically effective amount of a compound of Claim 1.

24. A method for the treatment of nasal congestion which comprises administering to a patient in need of such treatment a therapeutically effective amount of a compound of Claim 1.

25. A method for the treatment of allergic asthma which comprises administering to a patient in need of such treatment a therapeutically effective amount of a compound of Claim 1.

26. A method for the treatment of allergic rhinitis which comprises administering to a patient in need of such treatment a therapeutically effective amount of a compound of Claim 1.

27. A compound of formula I, as defined in any one of Claims 1 to 20, or a pharmaceutically acceptable salt or soluate thereof, for use in medicinal therapy.

5           28. A compound salt or hydrate of Claim 27 for use in treatment of prostaglandin D2 mediated diseases.

29. Use of a compound of formula I, as defined in any one of Claims 1 to 20, or a pharmaceutically acceptable salt or soluate thereof, in the  
10 manufacture of a medicament for treatment of nasal congestion, allergic asthma or allergic rhinitis.

30. A prostaglandin receptor antagonist pharmaceutical composition comprising an acceptable antagonistic amount of a compound of  
15 formula I, as defined in any one of Claims 1 to 20, or a pharmaceutically acceptable salt or soluate thereof, in association with a pharmaceutically acceptable carrier therefor.

## INTERNATIONAL SEARCH REPORT

International Classification No

PCT/CA 03/01658

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D471/14 A61K31/4375 A61P29/00 A61P11/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 128 364 A (GIRARD YVES ET AL) 7 July 1992 (1992-07-07) abstract; claims ---	1,21,22, 27
A	EP 0 441 517 A (PFIZER) 14 August 1991 (1991-08-14) claims 1-8 ---	1,21,27
A	EP 0 166 591 A (MERCK FROSST CANADA INC) 2 January 1986 (1986-01-02) claims 1,7-10 ---	1,21,22, 27
A	US 4 808 608 A (YOKIM CHRISTIANE ET AL) 28 February 1989 (1989-02-28) claims 1,10-12 --- -/--	1,21,22, 27



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the International filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

11 February 2004

Date of mailing of the international search report

23/02/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Hass, C



## INTERNATIONAL SEARCH REPORT

International Publication No

PCT/CA 03/01658

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 25919 A (HIRAMATSU YOSHIHARU; SHIONOGI & CO (JP); ARIMURA AKINORI (JP); HON) 18 June 1998 (1998-06-18) cited in the application claims -----	1,21-30
A	WO 99 62555 A (SHIONOGI & CO; ARIMURA AKINORI (JP)) 9 December 1999 (1999-12-09) cited in the application abstract -----	1,21-23, 27-30
A,P	WO 02 094830 A (MERCK FROSST CANADA INC; GUAY DANIEL (CA); DUFRESNE CLAUDE (CA); L) 28 November 2002 (2002-11-28) claims -----	1-30

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/CA 03/01658

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
  
Although claims 23-26 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/CA 03/01658

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5128364	A	07-07-1992	CA 2063955 A1	29-09-1992
EP 0441517	A	14-08-1991	AU 631723 B2	03-12-1992
			AU 7087991 A	29-08-1991
			BR 9100519 A	29-10-1991
			CA 2035805 C	27-06-1995
			CN 1054600 A	18-09-1991
			CS 9100299 A2	15-09-1991
			DE 9101348 U1	06-06-1991
			EG 19644 A	30-08-1995
			EP 0441517 A2	14-08-1991
			FI 910592 A	09-08-1991
			HU 70051 A2	28-09-1995
			IE 910402 A1	14-08-1991
			IL 97140 A	31-07-1995
			JP 4234845 A	24-08-1992
			JP 7119214 B	20-12-1995
			KR 9513562 B1	09-11-1995
			NO 910473 A ,B,	09-08-1991
			NZ 237040 A	26-10-1993
			PL 288977 A1	13-01-1992
			PL 167078 B1	31-07-1995
			PT 96681 A	31-10-1991
			US 5428043 A	27-06-1995
			ZA 9100916 A	30-09-1992
EP 0166591	A	02-01-1986	AU 4377685 A	02-01-1986
			CA 1241660 A1	06-09-1988
			CA 1288352 C2	03-09-1991
			DE 3574372 D1	28-12-1989
			DK 284485 A	24-02-1986
			EP 0166591 A2	02-01-1986
			ES 8609245 A1	16-12-1986
			GR 851494 A1	25-11-1985
			IL 75586 A	30-09-1988
			JP 61063653 A	01-04-1986
			PT 80698 A ,B	01-07-1985
			ZA 8504762 A	26-02-1986
US 4808608	A	28-02-1989	AU 611454 B2	13-06-1991
			AU 1926288 A	27-01-1989
			DK 404588 A	16-03-1989
			EP 0300676 A2	25-01-1989
			JP 1070462 A	15-03-1989
			NZ 225361 A	25-06-1991
			PT 88044 A ,B	30-06-1989
			US 5021447 A	04-06-1991
			ZA 8805209 A	29-03-1989
			CA 1305485 C	21-07-1992
			AT 61047 T	15-03-1991
			AU 592615 B2	18-01-1990
			AU 6790887 A	30-07-1987
			DE 3768097 D1	04-04-1991
			DK 34287 A	24-07-1987
			EP 0234708 A1	02-09-1987
			ES 2038653 T3	01-08-1993
			GR 3001828 T3	23-11-1992
			IE 59861 B1	20-04-1994

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Publication No

PCT/CA 03/01658

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4808608	A	IL 81325 A	29-11-1990
		JP 62223169 A	01-10-1987
		NZ 218999 A	21-12-1990
		PT 84178 A ,B	01-02-1987
		ZA 8700468 A	26-08-1987
WO 9825919	A 18-06-1998	AT 223911 T	15-09-2002
		AU 718958 B2	04-05-2000
		AU 5409798 A	03-07-1998
		BR 9714016 A	29-02-2000
		CN 1445225 A	01-10-2003
		CN 1240438 A ,B	05-01-2000
		CZ 9902065 A3	17-11-1999
		DE 69715448 D1	17-10-2002
		DE 69715448 T2	07-08-2003
		DK 944614 T3	13-01-2003
		EP 0944614 A1	29-09-1999
		ES 2181039 T3	16-02-2003
		HK 1020045 A1	13-12-2002
		HU 0002869 A2	28-06-2001
		ID 22753 A	09-12-1999
		IL 130660 A	10-03-2002
		IL 145863 A	24-06-2003
		IL 145864 A	24-06-2003
		IL 145865 A	31-07-2003
		WO 9825919 A1	18-06-1998
		JP 3215441 B2	09-10-2001
		JP 2000514824 T	07-11-2000
		NO 992838 A	20-07-1999
		NZ 336143 A	31-05-2002
		PL 334072 A1	31-01-2000
		PT 944614 T	31-01-2003
		RU 2161617 C1	10-01-2001
		TR 9901280 T2	21-10-1999
		TW 438790 B	07-06-2001
		US 6083974 A	04-07-2000
		ZA 9711039 A	15-06-1998
WO 9962555	A 09-12-1999	AU 3955199 A	20-12-1999
		CA 2333868 A1	09-12-1999
		CN 1304320 T	18-07-2001
		EP 1084711 A1	21-03-2001
		WO 9962555 A1	09-12-1999
		US 2003027854 A1	06-02-2003
		US 2002058693 A1	16-05-2002
WO 02094830	A 28-11-2002	WO 02094830 A2	28-11-2002